

COVID-19 10 STEP GUIDANCE FOR PHARMACY TEAMS

Government of Pakistan Ministry of National Health Services, Regulation & Coordination

1 Pharmacy Signage

Have a banner/standee at the pharmacy entrance advising patients not to enter the pharmacy if they are displaying any signs or symptoms of COVID-19. Signpost patients to contact the COVID-19 Helpline 1166 or to contact COVID-19 designated hospitals



Wash your hands

Regularly wash your hands with soap and water for at least 20 seconds or use an alcohol based rub. Use WHO 7 steps of hand-washing technique. Provide Hand gels at the pharmacy counter for the public



Wear a mask when in contact with patients.Change masks frequently. Disposable masks should only be used once.



Cash/PC Handling

Cash should be strictly handled with gloves and should be changed after every hour. Handling of medicines and cash should not be done by the same member of staff simultaneously. Use gloves to operate keyboards

Please



Self-Isolate

If you have a new cough and/ or a fever DO NOT come to work and selfisolate for at least 14 days and when symptoms get better



Wear disposable gloves in the pharmacy. Ensure you change your gloves every time you handle a new prescription



Sanitation of Premises

All surfaces including appliances, shelving, medicines packaging, computers, telephones should be sanitized on a regular basis using a rota system


🛛 📞 helpline 1166 If you are displaying symptoms



Maintain a 1 meter distance between yourself and patients when taking in and giving out prescriptions.Consider restricting the number of patients who can enter your pharmacy at one time



Encourage patients to sanitize mobile phones with alcohol wipes available at the pharmacy counter, before you handle the mobile



COVID-19 Testing

Testing will only take place if > Recently traveled internationally > Recently traveled Intercity and are showing COVID-19 Symptoms > Been in-contact with someone who has recently traveled Internationally >Anyone displaying COVID-19 symptoms: fever, persistent cough & Shortness of breadth



