

(5)  
REGISTERED NO. M - 302  
L - 7646



EXTRAORDINARY  
PUBLISHED BY AUTHORITY

---

ISLAMABAD, MONDAY, OCTOBER 16, 2023

---

PART II

Statutory Notifications (S. R. O.)

GOVERNMENT OF PAKISTAN  
MINISTRY OF NATIONAL HEALTH SERVICES,  
REGULATIONS & COORDINATION  
(Drug Regulatory Authority of Pakistan)

NOTIFICATION

*Islamabad, the 2nd October, 2023*

S. R. O. 1423(I)/2023.—For the purpose of clause (xi) of sub-paragraph (1) of paragraph 2 of the Drug Pricing Policy, 2018 the Ministry of National Health Services, Regulations & Coordination is hereby pleased to notify the Model List of Essential Medicines 23rd List (2023) published by the World Health Organization:

3181(1—69)

Price: Rs. 100.00

[1958(2023)/Ex.Gaz.]

---

**Explanatory notes**

The **core list** presents a list of minimum medicine needs for a basic health-care system, listing the most efficacious, safe and cost-effective medicines for priority conditions. Priority conditions are selected on the basis of current and estimated future public health relevance, and potential for safe and cost-effective treatment.

Where the [c] symbol is placed next to an individual medicine or strength of medicine on the core list it signifies that there is a specific indication for restricting its use to children.

The **complementary list** presents essential medicines for priority diseases, for which specialized diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training are needed. In case of doubt medicines may also be listed as complementary on the basis of consistent higher costs or less attractive cost-effectiveness in a variety of settings.

Where the [c/] symbol is placed next to an individual medicine or strength of medicine on the complementary list it signifies that the medicine(s) require(s) specialist diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training for their use in children.

The **square box symbol (□)** is intended to indicate therapeutic alternatives to the listed medicine that may be considered for selection in national essential medicines lists. Alternatives may be individual medicines, or multiple medicines within a pharmacological class or chemical subgroup, defined at the 4th level of the Anatomical Therapeutic Chemical (ATC) classification, which have similar clinical effectiveness and safety. The listed medicine should be the example of the class or subgroup for which there is the best evidence for effectiveness and safety. In some cases, this may be the first medicine that is licensed for marketing; in other instances, subsequently licensed compounds may be safer or more effective. Where there is no difference in terms of efficacy and safety data, the listed medicine should be the one that is generally available at the lowest price, based on international drug price information sources. Not all square box listings are applicable to medicine selection for children. A square box is not used to indicate alternative generic brands of the same small molecule medicines, nor alternative biosimilars of biological medicines. However, the selection and use of quality-assured generics and biosimilars of essential medicines at country level is recommended.

National lists should not use a similar symbol and should be specific in their final selection, which would depend on local availability and price.

The a symbol indicates that there is an age or weight restriction on use of the medicine; details for each medicine can be found in Table 1.1.

The presence of an entry on the Essential Medicines List carries no assurance as to pharmaceutical quality. It is the responsibility of the relevant national or regional drug regulatory authority to ensure that each product is of appropriate pharmaceutical quality (including stability) and that, when relevant, different products are interchangeable.

For recommendations and advice concerning all aspects of the quality assurance of medicines see the WHO website <https://www.who.int/teams/health-product-and-policy-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/guidelines/quality-assurance>

Medicines and dosage forms are listed in alphabetical order within each section and the order of listing does not imply preference for one form over another. Standard treatment guidelines should be consulted for information on appropriate dosage forms.

The main terms used for dosage forms in the Essential Medicines List can be found in Table 1.2.

Definitions of many of these terms and pharmaceutical quality requirements applicable to the different categories are published in the current edition of *The International Pharmacopoeia*. <https://www.who.int/teams/health-product-and-policy-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/pharmacopoeia>.

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>1. ANAESTHETICS, PREOPERATIVE MEDICINES AND MEDICAL GASES</b>	
<b>1.1 General anaesthetics and oxygen</b>	
<b>1.1.1 Inhalational medicines</b>	
halothane	Inhalation.
isoflurane	Inhalation.
nitrous oxide	Inhalation.
oxygen	Inhalation (medical gas).
sevoflurane	Inhalation.
<b>1.1.2 Injectable medicines</b>	
ketamine	Injection: 50 mg/mL (as hydrochloride) in 10 mL vial.
<input type="checkbox"/> propofol Therapeutic alternatives: - thiopental	Injection: 10 mg/mL; 20 mg/mL.
<b>1.2 Local anaesthetics</b>	
<input type="checkbox"/> bupivacaine Therapeutic alternatives to be reviewed	Injection: 0.25%; 0.5% (hydrochloride) in vial. Injection for spinal anaesthesia: 0.5% (hydrochloride) in 4 mL ampoule to be mixed with 7.5% glucose solution.
<input type="checkbox"/> lidocaine Therapeutic alternatives to be reviewed	Injection: 1%; 2% (hydrochloride) in vial. Injection for spinal anaesthesia: 5% (hydrochloride) in 2 mL ampoule to be mixed with 7.5% glucose solution. Topical forms: 2% to 4% (hydrochloride).
lidocaine + epinephrine (adrenaline)	Dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000. Injection: 1%; 2% (hydrochloride or sulfate) + epinephrine 1:200 000 in vial.
<i>Complementary List</i>	
ephedrine	Injection: 30 mg/mL (hydrochloride) in 1 mL ampoule. (For use in spinal anaesthesia during delivery, to prevent hypotension).
<b>1.3 Preoperative medication and sedation for short-term procedures</b>	
atropine	Injection: 1 mg (sulfate) in 1 mL ampoule.
<input type="checkbox"/> midazolam Therapeutic alternatives to be reviewed	Injection: 1 mg/mL. Oral liquid: 2 mg/mL [c]. Tablet: 7.5 mg; 15 mg.
morphine	Injection: 10 mg (sulfate or hydrochloride) in 1 mL ampoule.

**WHO Model List of Essential Medicines – 23rd List (2023)**

1.4 Medical gases	
oxygen*	<b>Inhalation</b> For use in the management of hypoxaemia. *No more than 30% oxygen should be used to initiate resuscitation of neonates less than or equal to 32 weeks of gestation.
2. MEDICINES FOR PAIN AND PALLIATIVE CARE	
2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)	
acetylsalicylic acid	<b>Suppository:</b> 50 mg to 150 mg. <b>Tablet:</b> 100 mg to 500 mg.
Ibuprofen <input checked="" type="checkbox"/>	<b>Oral liquid:</b> 100 mg/5 mL [c], 200 mg/5 mL. <b>Tablet:</b> 200 mg; 400 mg; 600 mg. <input checked="" type="checkbox"/> Not in children less than 3 months.
paracetamol (acetaminophen)*	<b>Oral liquid:</b> 120 mg/5 mL or 125 mg/5 mL**, 250 mg/5 mL [c]. **The presence of both 120 mg/5 mL and 125 mg/5 mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided. <b>Suppository:</b> 100 mg, 250 mg [c]. <b>Tablet:</b> 250 mg, 325 mg, 500 mg. <b>Tablet (dispersible):</b> 100 mg, 250 mg [c]. *Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.
2.2 Opioid analgesics	
codeine	<b>Tablet:</b> 30 mg (phosphate).
fentanyl*	<b>Transdermal patch:</b> 12 micrograms/hr; 25 micrograms/hr; 50 micrograms/hr; 75 micrograms/hr; 100 micrograms/hr. *For the management of cancer pain
<input checked="" type="checkbox"/> morphine Therapeutic alternatives: - hydromorphone - oxycodone	<b>Granules (slow release; to mix with water):</b> 20 mg to 200 mg (morphine sulfate). <b>Injection:</b> 10 mg (morphine hydrochloride or morphine sulfate) in 1 mL ampoule. <b>Oral liquid:</b> 10 mg/5 mL (morphine hydrochloride or morphine sulfate). <b>Tablet (slow release):</b> 10 mg to 200mg (morphine hydrochloride or morphine sulfate). <b>Tablet (immediate release):</b> 10 mg (morphine sulfate).
Complementary list	
methadone*	<b>Tablet:</b> 5 mg; 10 mg (hydrochloride) <b>Oral liquid:</b> 5 mg/5 mL; 10 mg/5 mL (hydrochloride) <b>Concentrate for oral liquid:</b> 5 mg/mL; 10 mg/mL (hydrochloride) *For the management of cancer pain.

### WHO Model List of Essential Medicines – 23rd List (2023)

2.3 Medicines for other common symptoms in palliative care	
amitriptyline	Tablet: 10 mg; 25 mg; 75 mg.  Injection: 50 mg/mL.
cyclizine [c]	Tablet: 50 mg.  Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule.
dexamethasone	Oral liquid: 2 mg/5 mL.  Tablet: 2 mg [c]; 4 mg.  Injection: 5 mg/mL.
diazepam	Oral liquid: 2 mg/5 mL.  Rectal gel: 5 mg/mL in 0.5 mL, 2 mL, 4 mL rectal delivery system.  Rectal solution: 2 mg/mL in 1.25 mL, 2.5 mL rectal tube; 4 mg/mL in 2.5 mL rectal tube.  Tablet: 5 mg; 10 mg.
docosate sodium	Capsule: 100 mg.  Oral liquid: 50 mg/5 mL.
fluoxetine	Solid oral dosage form: 20 mg (as hydrochloride).
haloperidol	Injection: 5 mg in 1 mL ampoule.  Oral liquid: 2 mg/mL.  Solid oral dosage form: 0.5 mg; 2mg; 5 mg.
hyoscine butylbromide	Injection: 20 mg/mL.
hyoscine hydrobromide [c]	Injection: 400 micrograms/mL; 600 micrograms/mL.  Transdermal patches: 1 mg/72 hours.
lactulose [c]	Oral liquid: 3.1 to 3.7 g/5 mL.
loperamide	Solid oral dosage form: 2 mg.
metoclopramide	Injection: 5 mg/mL (hydrochloride) in 2 mL ampoule.  Oral liquid: 5 mg/5 mL.  Solid oral form: 10 mg (hydrochloride).
midazolam	Injection: 1 mg/mL; 5 mg/mL.  Oral liquid: 2mg/mL [c].  Solid oral dosage form: 7.5 mg; 15 mg.
<input checked="" type="checkbox"/> ondansetron 	Therapeutic alternatives: - dolasetron - granisetron - palonosetron - tropisetron  Injection: 2 mg base/mL in 2 mL ampoule (as hydrochloride).  Oral liquid: 4 mg base/5 mL.  Solid oral dosage form: Eq 4 mg base; Eq 8 mg base.   > 1 month.
senna	Oral liquid: 7.5 mg/5 mL.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>3. ANTIALLERGICS AND MEDICINES USED IN ANAPHYLAXIS</b>	
dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule.
epinephrine (adrenaline)	Injection: 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
hydrocortisone	Powder for injection: 100 mg (as sodium succinate) in vial.
<input checked="" type="checkbox"/> loratadine*	Oral liquid: 1 mg/mL. Tablet: 10 mg.
Therapeutic alternatives: - cetirizine - fexofenadine	*There may be a role for sedating antihistamines for limited indications (EMLc).
<input checked="" type="checkbox"/> prednisolone	Oral liquid: 5 mg/mL [c]. Tablet: 5 mg; 25 mg.
Therapeutic alternatives: - prednisone	
<b>4. ANTIDOTES AND OTHER SUBSTANCES USED IN POISONINGS</b>	
<b>4.1 Non-specific</b>	
charcoal, activated	Powder.
<b>4.2 Specific</b>	
acetylcysteine	Injection: 200 mg/mL in 10 mL ampoule. Oral liquid: 10% [c]; 20% [o].
atropine	Injection: 1 mg (sulfate) in 1 mL ampoule.
calcium gluconate	Injection: 100 mg/mL in 10 mL ampoule.
methylthioninium chloride (methylene blue)	Injection: 10 mg/mL in 10 mL ampoule.
naloxone	Injection: 400 micrograms (hydrochloride) in 1 mL ampoule.
penicillamine	Solid oral dosage form: 250 mg.
potassium ferric hexacyano-ferrate(II) -2H <sub>2</sub> O (Prussian blue)	Powder for oral administration.
sodium nitrite	Injection: 30 mg/mL in 10 mL ampoule.
sodium thiosulfate	Injection: 250 mg/mL in 50 mL ampoule.
<b>Complementary List</b>	
deferoxamine	Powder for injection: 500 mg (mesilate) in vial.
dimercaprol	Injection in oil: 50 mg/mL in 2 mL ampoule.
fomepizole	Injection: 5 mg/mL (sulfate) in 20 mL ampoule or 1 g/mL (base) in 1.5 mL ampoule.
sodium calcium edetate	Injection: 200 mg/mL in 5 mL ampoule.
succimer	Solid oral dosage form: 100 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

### 5. MEDICINES FOR DISEASES OF THE NERVOUS SYSTEM

#### 5.1 Antiseizure medicines

carbamazepine	Oral liquid: 100 mg/5 mL. Tablet (chewable): 100 mg; 200 mg. Tablet (scored): 100 mg; 200 mg; 400 mg.
diazepam	Rectal gel: 5 mg/mL in 0.5 mL, 2 mL, 4 mL rectal delivery system. Rectal solution: 2 mg/mL in 1.25 mL, 2.5 mL rectal tube; 4 mg/mL in 2.5 mL rectal tube.
lamotrigine*	Tablet: 25 mg; 50 mg; 100 mg; 200 mg. Tablet (chewable, dispersible): 2 mg; 5 mg; 25 mg; 50 mg; 100 mg; 200 mg. *For use as adjunctive therapy for treatment-resistant partial or generalized seizures.
levetiracetam	Oral solution: 100 mg/mL. Tablet: 250 mg; 500 mg; 750 mg; 1000 mg.
<input checked="" type="checkbox"/> lorazepam Therapeutic alternatives: - diazepam (injection) - midazolam (injection)	Injection: 2 mg/mL in 1 mL ampoule; 4 mg/mL in 1 mL ampoule.
magnesium sulfate*	Injection: 0.5 g/mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume); 0.5 g/mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume). *For use in eclampsia and severe pre-eclampsia and not for other convulsant disorders.
midazolam	Solution for oromucosal administration: 5 mg/mL in 0.5 mL, 1 mL, 1.5 mL, 2 mL pre-filled syringe; 10 mg/mL in 0.25 mL, 0.5 mL, 0.75 mL, 1 mL pre-filled syringe. Injection*: 1 mg/mL in 5 mL vial; 5 mg/mL in 1 mL or 3 mL vial. *For buccal administration when solution for oromucosal administration is not available.
phenobarbital	Injection: 30 mg/mL or 60 mg/mL [c], 200 mg/mL (sodium). Oral liquid: 15 mg/5 mL. Tablet: 15 mg to 100 mg.
phenytoin	Injection: 50 mg/mL (phenytoin sodium). Oral liquid: 30 mg/5 mL (phenytoin). Solid oral dosage form: 25 mg; 50 mg; 100 mg (phenytoin sodium). Tablet (chewable): 50 mg (phenytoin).
valproic acid (sodium valproate)* <i>*Avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.</i>	Oral liquid: 200 mg/5 mL. Tablet (crushable): 100 mg. Tablet (enteric-coated): 200 mg; 500 mg.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>Complementary List</b>	
<i>ethosuximide</i>	<i>Capsule: 250 mg.</i> <i>Oral liquid: 250 mg/5 mL.</i>
<i>levetiracetam</i>	<i>Concentrate solution for infusion: 500 mg/5mL in 5 mL vial.</i> <i>Solution for infusion: 5 mg/mL; 10 mg/mL; 15 mg/mL in 100 mL bag.</i>
<i>valproic acid (sodium valproate)*</i>	<i>Injection: 100 mg/mL in 3 mL, 4 mL, 10 mL ampoule.</i>
<p>*<i>Avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.</i></p>	
<b>5.2 Medicines for multiple sclerosis</b>	
<b>Complementary List</b>	
<i>cladribine</i>	<i>Tablet: 10 mg.</i>
<i>glatiramer acetate</i>	<i>Injection (subcutaneous): 20 mg/mL; 40 mg/mL in pre-filled syringe.</i>
<i>rituximab*</i> *including quality-assured biosimilars	<i>Injection (intravenous): 500 mg/50 mL in 50 mL vial.</i>
<b>5.3 Medicines for parkinsonism</b>	
<input type="checkbox"/> <i>biperiden</i> Therapeutic alternatives: – trihexyphenidyl	<i>Injection: 5 mg (lactate) in 1 mL ampoule.</i> <i>Tablet: 2 mg (hydrochloride).</i>
<input type="checkbox"/> <i>levodopa</i> + <input type="checkbox"/> <i>carbidopa</i> Therapeutic alternatives: – benserazide (for carbidopa)	<i>Tablet: 100 mg + 10 mg; 100 mg + 25 mg; 250 mg + 25 mg.</i>

**WHO Model List of Essential Medicines – 23rd List (2023)****6. ANTI-INFECTIVE MEDICINES****6.1 Anthelmintics****6.1.1 Intestinal anthelmintics**

albendazole	Tablet (chewable, scored): 400 mg.
ivermectin	Tablet: 3 mg.
levamisole	Tablet: 50 mg; 150 mg (as hydrochloride).
mebendazole	Tablet (chewable): 100 mg; 500 mg.
niclosamide	Tablet (chewable): 500 mg.
praziquantel	Tablet: 150 mg, 500 mg Tablet (scored): 600 mg.
pyrantel	Tablet (chewable): 250 mg (as embonate or pamoate).

**6.1.2 Antifilarials**

albendazole	Tablet (chewable, scored): 400 mg.
diethylcarbamazine	Tablet: 50 mg; 100 mg (dihydrogen citrate).
ivermectin	Tablet: 3 mg.

**6.1.3 Antischistosomals and other antitrematode medicines**

praziquantel	Tablet: 150 mg, 500 mg. Tablet (scored): 600 mg.
triclabendazole	Tablet (scored): 250 mg.

**Complementary List**

oxamniquine*	Capsule: 250 mg. Oral liquid: 250 mg/5 mL.
*For use when praziquantel treatment fails.	

**6.1.4 Cysticidal medicines****Complementary List**

albendazole	Tablet (chewable): 200 mg [c]. Tablet (chewable, scored): 400 mg.
mebendazole	Tablet (chewable): 100 mg [c], 500 mg.
praziquantel	Tablet: 150 mg, 500 mg. Tablet (scored): 600 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

### 6.2 Antibacterials

To assist in the development of tools for antibiotic stewardship at local, national and global levels and to reduce antimicrobial resistance, the Access, Watch, Reserve (AWaRe) classification of antibiotics was developed – where antibiotics are classified into different groups to emphasize the importance of their appropriate use.

#### ACCESS GROUP ANTIBIOTICS

This group includes antibiotics that have activity against a wide range of commonly encountered susceptible pathogens while also showing lower resistance potential than antibiotics in the other groups. Selected Access group antibiotics are recommended as essential first or second choice empiric treatment options for infectious syndromes reviewed by the EML Expert Committee and are listed as individual medicines on the Model Lists to improve access and promote appropriate use. They are essential antibiotics that should be widely available, affordable and quality assured.

#### WATCH GROUP ANTIBIOTICS

This group includes antibiotic classes that have higher resistance potential and includes most of the highest priority agents among the Critically Important Antimicrobials for Human Medicine and/or antibiotics that are at relatively high risk of selection of bacterial resistance. These medicines should be prioritized as key targets of stewardship programs and monitoring. Selected Watch group antibiotics are recommended as essential first or second choice empiric treatment options for a limited number of specific infectious syndromes and are listed as individual medicines on the Model Lists.

#### RESERVE GROUP ANTIBIOTICS

This group includes antibiotics and antibiotic classes that should be reserved for treatment of confirmed or suspected infections due to multi-drug-resistant organisms. Reserve group antibiotics should be treated as "last resort" options. Selected Reserve group antibiotics are listed as individual medicines on the Model Lists when they have a favourable risk-benefit profile and proven activity against "Critical Priority" or "High Priority" pathogens identified by the WHO Priority Pathogens List, notably carbapenem resistant *Enterobacteriaceae*. These antibiotics should be accessible, but their use should be tailored to highly specific patients and settings, when all alternatives have failed or are not suitable. These medicines could be protected and prioritized as key targets of national and international stewardship programs involving monitoring and utilization reporting, to preserve their effectiveness.

## WHO Model List of Essential Medicines – 23rd List (2023)

6.2.1 Access group antibiotics						
amikacin	<p><b>Injection:</b> 50 mg/mL (as sulfate) [c]; 250 mg/mL (as sulfate) in 2 mL vial.</p> <table> <tr> <td><b>FIRST CHOICE</b></td><td><b>SECOND CHOICE</b></td></tr> <tr> <td> <ul style="list-style-type: none"> <li>– <i>High-risk febrile neutropenia</i></li> <li>– <i>Pyelonephritis or prostatitis (severe)</i></li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Sepsis in neonates and children</i> [c]</li> </ul> </td></tr> </table>	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<ul style="list-style-type: none"> <li>– <i>High-risk febrile neutropenia</i></li> <li>– <i>Pyelonephritis or prostatitis (severe)</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Sepsis in neonates and children</i> [c]</li> </ul>	
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<ul style="list-style-type: none"> <li>– <i>High-risk febrile neutropenia</i></li> <li>– <i>Pyelonephritis or prostatitis (severe)</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Sepsis in neonates and children</i> [c]</li> </ul>					
amoxicillin	<p><b>Powder for injection:</b> 250 mg; 500 mg; 1 g (as sodium) in vial.</p> <p><b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL (as trihydrate) [c].</p> <p><b>Solid oral dosage form:</b> 250 mg; 500 mg; 1g (as trihydrate).</p> <p><b>Tablet (dispersible, scored):</b> 250 mg; 500 mg (as trihydrate) [c].</p> <table> <tr> <td><b>FIRST CHOICE</b></td><td><b>SECOND CHOICE</b></td></tr> <tr> <td> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated severe acute malnutrition</i> [c]</li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Pharyngitis</i></li> <li>– <i>Progressive apical dental abscess</i></li> <li>– <i>Sepsis in neonates and children</i> [c]</li> <li>– <i>Sinusitis</i></li> <li>– <i>Uncomplicated severe acute malnutrition</i> [c]</li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul> </td></tr> </table>	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated severe acute malnutrition</i> [c]</li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Pharyngitis</i></li> <li>– <i>Progressive apical dental abscess</i></li> <li>– <i>Sepsis in neonates and children</i> [c]</li> <li>– <i>Sinusitis</i></li> <li>– <i>Uncomplicated severe acute malnutrition</i> [c]</li> </ul>	<ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul>	
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated severe acute malnutrition</i> [c]</li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Pharyngitis</i></li> <li>– <i>Progressive apical dental abscess</i></li> <li>– <i>Sepsis in neonates and children</i> [c]</li> <li>– <i>Sinusitis</i></li> <li>– <i>Uncomplicated severe acute malnutrition</i> [c]</li> </ul>	<ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul>					
amoxicillin + clavulanic acid	<p><b>Powder for injection:</b> 500 mg (as sodium) + 100 mg (as potassium salt); 1000 mg (as sodium) + 200 mg (as potassium salt) in vial.</p> <p><b>Powder for oral liquid:</b> 125 mg (as trihydrate)+ 31.25 mg (as potassium salt)/5 mL; 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5mL [c].</p> <p><b>Tablet:</b> 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt).</p> <p><b>Tablet (dispersible):</b> 200 mg (as trihydrate) + 28.5 mg (as potassium salt) [c]; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) [c].</p> <table> <tr> <td><b>FIRST CHOICE</b></td><td><b>SECOND CHOICE</b></td></tr> <tr> <td> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Hospital acquired pneumonia</i></li> <li>– <i>Low-risk febrile neutropenia</i></li> <li>– <i>Lower urinary tract infections</i></li> <li>– <i>Sinusitis</i></li> <li>– <i>Skin and soft tissue infections</i></li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Bone and joint infections</i></li> <li>– <i>Community-acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Surgical prophylaxis</i></li> </ul> </td></tr> </table>	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Hospital acquired pneumonia</i></li> <li>– <i>Low-risk febrile neutropenia</i></li> <li>– <i>Lower urinary tract infections</i></li> <li>– <i>Sinusitis</i></li> <li>– <i>Skin and soft tissue infections</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Bone and joint infections</i></li> <li>– <i>Community-acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Surgical prophylaxis</i></li> </ul>	
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Hospital acquired pneumonia</i></li> <li>– <i>Low-risk febrile neutropenia</i></li> <li>– <i>Lower urinary tract infections</i></li> <li>– <i>Sinusitis</i></li> <li>– <i>Skin and soft tissue infections</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Bone and joint infections</i></li> <li>– <i>Community-acquired pneumonia (mild to moderate)</i></li> <li>– <i>Community acquired pneumonia (severe)</i></li> <li>– <i>Otitis media</i></li> <li>– <i>Surgical prophylaxis</i></li> </ul>					

## WHO Model List of Essential Medicines – 23rd List (2023)

	<b>Powder for injection:</b> 500 mg; 1 g (as sodium) in vial.	
ampicillin	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Community acquired pneumonia (severe) [c]</li> <li>– Complicated intraabdominal infections [c]</li> <li>– Complicated severe acute malnutrition [c]</li> <li>– Sepsis in neonates and children [c]</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Acute bacterial meningitis</li> </ul>
benzathine benzylpenicillin	<b>Powder for injection:</b> 1.2 million IU (~ 900 mg) in vial [c]; 2.4 million IU (~ 1.8 g) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Syphilis</li> </ul>	<b>SECOND CHOICE</b>
benzylpenicillin	<b>Powder for injection:</b> 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Community acquired pneumonia (severe) [c]</li> <li>– Complicated severe acute malnutrition [c]</li> <li>– Sepsis in neonates and children [c]</li> <li>– Syphilis</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Acute bacterial meningitis</li> </ul>
cefalexin	<b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL (anhydrous). <b>Solid oral dosage form:</b> 250 mg; 500 mg (as monohydrate). <b>Tablet (dispersible):</b> 125 mg [c]; 250 mg [c].	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Skin and soft tissue infections</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Exacerbations of COPD</li> <li>– Pharyngitis</li> </ul>
cefazin 	<b>Powder for injection:</b> 1 g (as sodium salt) in vial. <input checked="" type="checkbox"/> > 1 month.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Surgical prophylaxis</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Bone and joint infections</li> </ul>
chloramphenicol	<b>Oily suspension for injection*</b> : 0.5 g/mL (as sodium succinate) in 2 mL ampoule. *Only for the presumptive treatment of epidemic meningitis in children older than 2 years and in adults. <b>Powder for injection:</b> 1 g (as sodium succinate) in vial.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Acute bacterial meningitis</li> </ul>

## WHO Model List of Essential Medicines – 23rd List (2023)

	<b>clindamycin</b>	<b>Capsule:</b> 150 mg (as hydrochloride). <b>Injection:</b> 150 mg/mL (as phosphate); 600 mg/4 mL (as phosphate); 900 mg/6 mL (as phosphate). <b>Powder for oral liquid:</b> 75 mg/5 mL (as palmitate hydrochloride) [c].	
		<b>FIRST CHOICE</b> – <i>Necrotizing fasciitis</i>	<b>SECOND CHOICE</b> – <i>Bone and joint infections</i>
	<b>□ cloxacillin*</b> Therapeutic alternatives: – 4 <sup>th</sup> level ATC chemical subgroup (J01CF Beta-lactamase resistant penicillins)	<b>Capsule:</b> 250 mg [c], 500 mg; 1 g (as sodium). <b>Powder for injection:</b> 250 mg [c], 500 mg (as sodium) in vial. <b>Powder for oral liquid:</b> 125 mg/5 mL, 250 mg/5 mL (as sodium) [c]. *cloxacillin, dicloxacillin and flucloxacillin are preferred for oral administration due to better bioavailability.	
		<b>FIRST CHOICE</b> – <i>Bone and joint infections</i> – <i>Skin and soft tissue infections</i>	<b>SECOND CHOICE</b> – <i>Sepsis in neonates and children</i> [c]
	<b>doxycycline □</b>	<b>Oral liquid:</b> 50 mg/5 mL (calcium) [c]. <b>Powder for oral liquid:</b> 25 mg/5 mL (monohydrate) [c]. <b>Powder for injection:</b> 100 mg in vial. <b>Solid oral dosage form:</b> 50 mg [c]; 100 mg (as hydiate). <b>Tablet (dispersible):</b> 100 mg (as monohydrate) [c].	
		[c] Use in children <8 years only for life-threatening infections when no alternative exists.	
	<b>gentamicin</b>	<b>Injection:</b> 10 mg/mL (as sulfate); 40 mg/mL (as sulfate) in 2 mL vial.	
		<b>FIRST CHOICE</b> – <i>Acute bacterial meningitis in neonates</i> [c] – <i>Community acquired pneumonia (severe)</i> [c] – <i>Complicated intraabdominal infections</i> [c] – <i>Complicated severe acute malnutrition</i> [c] – <i>Sepsis in neonates and children</i> [c]	<b>SECOND CHOICE</b> – <i>Gonorrhoea</i> – <i>Surgical prophylaxis</i>

## WHO Model List of Essential Medicines – 23rd List (2023)

	<p><b>Injection:</b> 500 mg in 100 mL vial.  <b>Oral liquid:</b> 200 mg/5 mL (as benzoate).  <b>Suppository:</b> 500 mg; 1 g.  <b>Tablet:</b> 200 mg; 250 mg; 400 mg; 500 mg.</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>– <i>C. difficile infection</i></li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Complicated Intrabdominal infections (severe)</i></li> <li>– <i>Necrotizing fascitis</i></li> <li>– <i>Surgical prophylaxis</i></li> <li>– <i>Trichomoniasis</i></li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> </ul> </td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE	<ul style="list-style-type: none"> <li>– <i>C. difficile infection</i></li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Complicated Intrabdominal infections (severe)</i></li> <li>– <i>Necrotizing fascitis</i></li> <li>– <i>Surgical prophylaxis</i></li> <li>– <i>Trichomoniasis</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> </ul>
FIRST CHOICE	SECOND CHOICE				
<ul style="list-style-type: none"> <li>– <i>C. difficile infection</i></li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Complicated Intrabdominal infections (severe)</i></li> <li>– <i>Necrotizing fascitis</i></li> <li>– <i>Surgical prophylaxis</i></li> <li>– <i>Trichomoniasis</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> </ul>				
metronidazole	<p><b>Oral liquid:</b> 25 mg/5 mL [c].  <b>Solid oral dosage form:</b> 50 mg [c]; 100 mg.</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul> </td><td></td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE	<ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul>	
FIRST CHOICE	SECOND CHOICE				
<ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul>					
nitrofurantoin	<p><b>Powder for oral liquid:</b> 250 mg/5 mL (as potassium).  <b>Solid oral dosage form:</b> 250 mg; 500 mg (as potassium).</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td></td><td></td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE		
FIRST CHOICE	SECOND CHOICE				
phenoxymethylenicillin	<p><b>FIRST CHOICE</b></p> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (mild to moderate)</i></li> <li>– <i>Pharyngitis</i></li> <li>– <i>Progressive apical dental abscess</i></li> </ul> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td></td><td></td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE		
FIRST CHOICE	SECOND CHOICE				
procaine benzylpenicillin*	<p><b>Powder for injection:</b> 1 g (=1 million IU); 3 g (=3 million IU) in vial.</p> <p>*Procaine benzylpenicillin is not recommended as first-line treatment for neonatal sepsis except in settings with high neonatal mortality, when given by trained health workers in cases where hospital care is not achievable.</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>– <i>Syphilis (congenital)</i> [c]</li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Syphilis</i></li> </ul> </td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE	<ul style="list-style-type: none"> <li>– <i>Syphilis (congenital)</i> [c]</li> </ul>	<ul style="list-style-type: none"> <li>– <i>Syphilis</i></li> </ul>
FIRST CHOICE	SECOND CHOICE				
<ul style="list-style-type: none"> <li>– <i>Syphilis (congenital)</i> [c]</li> </ul>	<ul style="list-style-type: none"> <li>– <i>Syphilis</i></li> </ul>				
spectinomycin	<p><b>Powder for injection:</b> 2 g (as hydrochloride) in vial.</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td></td><td> <ul style="list-style-type: none"> <li>– <i>Gonorrhoea</i></li> </ul> </td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE		<ul style="list-style-type: none"> <li>– <i>Gonorrhoea</i></li> </ul>
FIRST CHOICE	SECOND CHOICE				
	<ul style="list-style-type: none"> <li>– <i>Gonorrhoea</i></li> </ul>				
sulfamethoxazole + trimethoprim	<p><b>Injection:</b> 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule.</p> <p><b>Oral liquid:</b> 200 mg + 40 mg/5 mL.</p> <p><b>Tablet:</b> 100 mg + 20 mg; 400 mg + 80 mg; 800 mg + 160 mg.</p> <p><b>Tablet (dispersible):</b> 100 mg + 20 mg [c].</p> <table border="1"> <thead> <tr> <th>FIRST CHOICE</th><th>SECOND CHOICE</th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul> </td><td> <ul style="list-style-type: none"> <li>– <i>Acute invasive diarrhoea / bacterial dysentery</i></li> </ul> </td></tr> </tbody> </table>	FIRST CHOICE	SECOND CHOICE	<ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Acute invasive diarrhoea / bacterial dysentery</i></li> </ul>
FIRST CHOICE	SECOND CHOICE				
<ul style="list-style-type: none"> <li>– <i>Lower urinary tract infections</i></li> </ul>	<ul style="list-style-type: none"> <li>– <i>Acute invasive diarrhoea / bacterial dysentery</i></li> </ul>				

## WHO Model List of Essential Medicines – 23rd List (2023)

trimethoprim	Tablet: 100 mg; 200 mg. Oral liquid: 50 mg/5 mL [c].	
	FIRST CHOICE <ul style="list-style-type: none"><li>– Lower urinary tract infections</li></ul>	SECOND CHOICE
<b>6.2.2 Watch group antibiotics</b>		
azithromycin	Solid oral dosage form: 250 mg; 500 mg (anhydrous). Powder for oral liquid: 200 mg/5 mL (anhydrous) [c].	
	FIRST CHOICE <ul style="list-style-type: none"><li>– Cholera</li><li>– Enteric fever</li><li>– Gonorrhoea</li><li>– Sexually transmitted infection due to <i>Chlamydia trachomatis</i></li><li>– Trachoma</li><li>– Yaws</li></ul>	SECOND CHOICE <ul style="list-style-type: none"><li>– Acute invasive bacterial diarrhoea / dysentery</li><li>– Gonorrhoea</li></ul>
cefixime	Powder for oral liquid: 100 mg/5 mL [c]. Solid oral dosage form: 200 mg; 400 mg (as trihydrate).	
	FIRST CHOICE	SECOND CHOICE <ul style="list-style-type: none"><li>– Acute invasive bacterial diarrhoea / dysentery</li><li>– Gonorrhoea</li></ul>
cefotaxime*	Powder for injection: 250 mg; 500 mg; 1 g; 2 g (as sodium) in vial. *3rd generation cephalosporin of choice for use in hospitalized neonates.	
	FIRST CHOICE <ul style="list-style-type: none"><li>– Acute bacterial meningitis</li><li>– Community acquired pneumonia (severe)</li><li>– Complicated intraabdominal infections (mild to moderate)</li><li>– Complicated intraabdominal infections (severe)</li><li>– Hospital acquired pneumonia</li><li>– Pyelonephritis or prostatitis (severe)</li></ul>	SECOND CHOICE <ul style="list-style-type: none"><li>– Bone and joint infections</li><li>– Pyelonephritis or prostatitis (mild to moderate)</li><li>– Sepsis in neonates and children [c]</li></ul>

## WHO Model List of Essential Medicines – 23rd List (2023)

	<p><b>Powder for Injection:</b> 250 mg; 500 mg; 1 g; 2 g (as sodium) in vial.        *Do not administer with calcium and avoid in infants with hyperbilirubinaemia.  <input checked="" type="checkbox"/> &gt; 41 weeks corrected gestational age.</p> <table border="1"> <thead> <tr> <th style="text-align: center;"><b>FIRST CHOICE</b></th><th style="text-align: center;"><b>SECOND CHOICE</b></th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>- Acute bacterial meningitis</li> <li>- Community acquired pneumonia (severe)</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> <li>- Complicated intrabdominal infections (severe)</li> <li>- Endophthalmitis</li> <li>- Enteric fever</li> <li>- Gonorrhoea</li> <li>- Hospital acquired pneumonia</li> <li>- Necrotizing fasciitis</li> <li>- Pyelonephritis or prostatitis (severe)</li> </ul> </td><td> <ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Bone and joint infections</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> <li>- Sepsis in neonates and children [c]</li> </ul> </td></tr> </tbody> </table>		<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<ul style="list-style-type: none"> <li>- Acute bacterial meningitis</li> <li>- Community acquired pneumonia (severe)</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> <li>- Complicated intrabdominal infections (severe)</li> <li>- Endophthalmitis</li> <li>- Enteric fever</li> <li>- Gonorrhoea</li> <li>- Hospital acquired pneumonia</li> <li>- Necrotizing fasciitis</li> <li>- Pyelonephritis or prostatitis (severe)</li> </ul>	<ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Bone and joint infections</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> <li>- Sepsis in neonates and children [c]</li> </ul>
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<ul style="list-style-type: none"> <li>- Acute bacterial meningitis</li> <li>- Community acquired pneumonia (severe)</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> <li>- Complicated intrabdominal infections (severe)</li> <li>- Endophthalmitis</li> <li>- Enteric fever</li> <li>- Gonorrhoea</li> <li>- Hospital acquired pneumonia</li> <li>- Necrotizing fasciitis</li> <li>- Pyelonephritis or prostatitis (severe)</li> </ul>	<ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Bone and joint infections</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> <li>- Sepsis in neonates and children [c]</li> </ul>					
ceftriaxone* <input checked="" type="checkbox"/>	<p><b>Powder for Injection:</b> 250 mg; 750 mg; 1.5 g (as sodium) in vial.</p> <table border="1"> <thead> <tr> <th style="text-align: center;"><b>FIRST CHOICE</b></th><th style="text-align: center;"><b>SECOND CHOICE</b></th></tr> </thead> <tbody> <tr> <td></td><td> <ul style="list-style-type: none"> <li>- Surgical prophylaxis</li> </ul> </td></tr> </tbody> </table>		<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>		<ul style="list-style-type: none"> <li>- Surgical prophylaxis</li> </ul>
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
	<ul style="list-style-type: none"> <li>- Surgical prophylaxis</li> </ul>					
ciprofloxacin	<p><b>Oral liquid:</b> 250 mg/5 mL (anhydrous) [c].  <b>Solution for IV Infusion:</b> 2 mg/mL (as hydrate) [c].  <b>Solid oral dosage form:</b> 100 mg [c]; 250 mg; 500 mg (as hydrochloride).</p> <table border="1"> <thead> <tr> <th style="text-align: center;"><b>FIRST CHOICE</b></th><th style="text-align: center;"><b>SECOND CHOICE</b></th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Enteric fever</li> <li>- Low-risk febrile neutropenia</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> </ul> </td><td> <ul style="list-style-type: none"> <li>- Cholera</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> </ul> </td></tr> </tbody> </table>		<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Enteric fever</li> <li>- Low-risk febrile neutropenia</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> </ul>	<ul style="list-style-type: none"> <li>- Cholera</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> </ul>
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<ul style="list-style-type: none"> <li>- Acute invasive bacterial diarrhoea / dysentery</li> <li>- Enteric fever</li> <li>- Low-risk febrile neutropenia</li> <li>- Pyelonephritis or prostatitis (mild to moderate)</li> </ul>	<ul style="list-style-type: none"> <li>- Cholera</li> <li>- Complicated intraabdominal infections (mild to moderate)</li> </ul>					
<input checked="" type="checkbox"/> clarithromycin† Therapeutic alternatives: - erythromycin*	<p><b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL.  <b>Powder for Injection:</b> 500 mg in vial.  <b>Solid oral dosage form:</b> 250 mg [c]; 500 mg.</p> <p>†clarithromycin is also listed for use in combination regimens for eradication of <i>H. pylori</i> in adults.</p> <table border="1"> <thead> <tr> <th style="text-align: center;"><b>FIRST CHOICE</b></th><th style="text-align: center;"><b>SECOND CHOICE</b></th></tr> </thead> <tbody> <tr> <td><i>Community acquired pneumonia (severe)</i></td><td><i>- Pharyngitis</i></td></tr> </tbody> </table>		<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>	<i>Community acquired pneumonia (severe)</i>	<i>- Pharyngitis</i>
<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b>					
<i>Community acquired pneumonia (severe)</i>	<i>- Pharyngitis</i>					

**WHO Model List of Essential Medicines – 23rd List (2023)**

piperacillin + tazobactam	<b>Powder for injection:</b> 2 g (as sodium) + 250 mg (as sodium); 4 g (as sodium) + 500 mg (as sodium) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Complicated intraabdominal infections (severe)</i></li> <li>– <i>High-risk febrile neutropenia</i></li> <li>– <i>Hospital acquired pneumonia</i></li> <li>– <i>Necrotizing fasciitis</i></li> </ul>	<b>SECOND CHOICE</b>
vancomycin*	<b>Capsule:</b> 125 mg; 250 mg (as hydrochloride). *vancomycin powder for injection may also be used for oral administration	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>C. difficile infection</i></li> </ul>
<b>Complementary List</b>		
ceftazidime	<b>Powder for injection:</b> 250 mg; 1 g (as pentahydrate) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Endophthalmitis</i></li> </ul>	<b>SECOND CHOICE</b>
<input checked="" type="checkbox"/> meropenem*  <i>Therapeutic alternatives:</i> - imipenem + cilastatin  *complicated intraabdominal infections and high-risk febrile neutropenia only. Meropenem is the preferred choice for acute bacterial meningitis in neonates.	<b>Powder for injection:</b> 500 mg (as trihydrate); 1 g (as trihydrate) in vial.  <input checked="" type="checkbox"/> > 3 months.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis in neonates [c]</i></li> <li>– <i>Complicated intraabdominal infections (severe)</i></li> <li>– <i>High-risk febrile neutropenia</i></li> </ul>
vancomycin	<b>Powder for injection:</b> 250 mg; 500 mg; 1 g (as hydrochloride) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Endophthalmitis</i></li> <li>– <i>Necrotizing fasciitis</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>High-risk febrile neutropenia</i></li> </ul>
<b>6.2.3 Reserve group antibiotics</b>		
<b>Complementary List</b>		
cefiderocol	<b>Powder for injection:</b> 1 g (as sulfate toxylate) in vial.	
ceftazidime + avibactam	<b>Powder for injection:</b> 2 g + 0.5 g in vial.	
ceftolozane + tazobactam	<b>Powder for injection:</b> 1 g + 0.5 g in vial.	
colistin	<b>Powder for injection:</b> 1 million IU (as colistimethate sodium) (equivalent to 34 mg colistin base activity) in vial.	
fosfomycin	<b>Powder for injection:</b> 2 g; 4 g (as sodium) in vial.	
<input checked="" type="checkbox"/> linezolid  <i>Therapeutic alternatives:</i> - tedizolid phosphate	<b>Injection for intravenous administration:</b> 2 mg/mL in 300 mL bag.  <b>Powder for oral liquid:</b> 100 mg/5 mL.  <b>Tablet:</b> 600 mg.  <b>Tablet (dispersible):</b> 150 mg [c].	

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>meropenem + vaborbactam</i>	<i>Powder for injection: 1 g (as trihydrate) + 1 g in vial.</i>
<i>plazomicin</i>	<i>Injection: 500 mg/10 mL.</i>
<i>polymyxin B</i>	<i>Powder for injection: 500 000 IU (equivalent to 50 mg polymyxin B base) in vial.</i>
<b>6.2.4 Antileprosy medicines</b>	
Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance. Colour-coded blister packs (MDT blister packs) containing standard two-medicine (paucibacillary leprosy) or three-medicine (multibacillary leprosy) combinations for adult and childhood leprosy should be used. MDT blister packs can be supplied free of charge through WHO.	
<i>clofazimine</i>	<i>Solid oral dosage form: 50 mg; 100 mg.</i>
<i>dapsone</i>	<i>Tablet: 25 mg; 50 mg; 100 mg.</i>
<i>rifampicin</i>	<i>Oral liquid: 20 mg/mL [c].</i> <i>Solid oral dosage form: 150 mg; 300 mg.</i>
<b>6.2.5 Antituberculosis medicines</b>	
WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.	
<i>ethambutol</i>	<i>Tablet: 100 mg; 400 mg (hydrochloride).</i> <i>Tablet (dispersible): 100 mg [c].</i>
<i>ethambutol + isoniazid + pyrazinamide + rifampicin</i>	<i>Tablet: 275 mg + 75 mg + 400 mg + 150 mg.</i>
<i>ethambutol + isoniazid + rifampicin</i>	<i>Tablet: 275 mg + 75 mg + 150 mg.</i>
<i>ethionamide</i>	<i>Tablet: 250 mg.</i> <i>Tablet (dispersible): 125 mg [c].</i>
<i>isoniazid</i>	<i>Tablet: 100 mg; 300 mg.</i> <i>Tablet (dispersible): 100 mg [c].</i>
<i>isoniazid + pyrazinamide + rifampicin</i>	<i>Tablet (dispersible): 50 mg + 150 mg + 75 mg [c].</i>
<i>isoniazid + rifampicin</i>	<i>Tablet: 75 mg + 150 mg; 150 mg + 300 mg.</i> <i>Tablet (dispersible): 50 mg + 75 mg [c].</i>
<i>isoniazid + rifapentine</i>	<i>Tablet (scored): 300 mg + 300 mg.</i>
<i>moxifloxacin</i>	<i>Tablet: 400 mg.</i>
<i>pyrazinamide</i>	<i>Tablet: 400 mg; 500 mg.</i> <i>Tablet (dispersible): 150 mg.</i>
<i>rifabutin</i>	<i>Solid oral dosage form: 150 mg.*</i> *For use only in patients with HIV receiving protease inhibitors.
<i>rifampicin</i>	<i>Oral liquid: 20 mg/mL [c].</i> <i>Solid oral dosage form: 150 mg; 300 mg.</i>
<i>rifapentine</i>	<i>Tablet: 150 mg; 300 mg.</i>

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>Complementary List</b>	
Medicines for the treatment of multidrug-resistant tuberculosis (MDR-TB) should be used in specialized centres adhering to WHO standards for TB control.	
amikacin	<b>Injection:</b> 250 mg/mL (as sulfate) in 2 mL vial.
amoxicillin + clavulanic acid*	<b>Powder for oral liquid:</b> 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5mL [c]. <b>Tablet:</b> 500 mg (as trihydrate) + 125 mg (as potassium salt). *For use only in combination with meropenem or imipenem+cilastatin.
bedaquiline	<b>Tablet:</b> 20 mg [c]; 100 mg.
clofazimine	<b>Solid oral dosage form:</b> 50 mg; 100 mg.
<input checked="" type="checkbox"/> cycloserine	<b>Solid oral dosage form:</b> 125 mg [c]; 250 mg.
Therapeutic alternatives: - terizidone	
delamanid	<b>Tablet (dispersible):</b> 25 mg [c]. <b>Tablet:</b> 50 mg.
<input checked="" type="checkbox"/> ethionamide	<b>Tablet:</b> 250 mg. <b>Tablet (dispersible):</b> 125 mg [c].
Therapeutic alternatives: - prontosilamide	
levofloxacin	<b>Tablet:</b> 250mg; 500 mg; 750 mg. <b>Tablet (dispersible):</b> 100 mg [c].
linezolid	<b>Tablet:</b> 600 mg. <b>Tablet (dispersible):</b> 150 mg [c].
<input checked="" type="checkbox"/> meropenem	<b>Powder for Injection:</b> 500 mg (as trihydrate); 1 g (as trihydrate) in vial. - imipenem + cilastatin
moxifloxacin	<b>Tablet:</b> 400 mg. <b>Tablet (dispersible):</b> 100 mg [c].
p-aminosalicylate sodium	<b>Powder for oral solution:</b> 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid).
pretomanid	<b>Tablet:</b> 200 mg.
streptomycin [c]	<b>Powder for injection:</b> 1 g (as sulfate) in vial.

## WHO Model List of Essential Medicines – 23rd List (2023)

6.3 Antifungal medicines	
amphotericin B*	<p>Powder for injection: 50 mg (liposomal complex) in vial.</p> <p>Powder for injection: 50 mg (as sodium deoxycholate) in vial</p> <p>*Liposomal amphotericin B has a better safety profile than the sodium deoxycholate formulation and should be prioritized for selection and use depending on local availability and cost.</p>
clotrimazole	<p>Vaginal cream: 1%; 10%.</p> <p>Vaginal tablet: 100 mg; 500 mg.</p>
fluconazole	<p>Capsule: 50 mg.</p> <p>Injection: 2 mg/ml. In vial.</p> <p>Oral liquid: 50 mg/5 mL.</p> <p>Powder for oral liquid: 50 mg/5 mL [c].</p>
flucytosine	<p>Capsule: 250 mg.</p> <p>Infusion: 2.5 g in 250 mL.</p>
griseofulvin	<p>Oral liquid: 125 mg/5 mL [c].</p> <p>Solid oral dosage form: 125 mg; 250 mg.</p>
itraconazole*	<p>Capsule: 100 mg.</p> <p>Oral liquid: 10 mg/mL.</p> <p>*For treatment of chronic pulmonary aspergillosis, histoplasmosis, sporotrichosis, paracoccidioidomycosis, mycoses caused by <i>T. martei</i> and chromoblastomycosis; and prophylaxis of histoplasmosis and infections caused by <i>T. martei</i> in AIDS patients.</p>
nystatin	<p>Lozenge: 100 000 IU.</p> <p>Oral liquid: 100 000 IU/mL [c].</p> <p>Pessary: 100 000 IU.</p> <p>Solid oral dosage form: 500 000 IU.</p>
voriconazole*	<p>Tablet: 50 mg; 200 mg</p> <p>Powder for injection: 200 mg in vial</p> <p>Powder for oral liquid: 40 mg/mL.</p> <p>*For treatment of chronic pulmonary aspergillosis and acute invasive aspergillosis.</p>
Complementary List	
<input checked="" type="checkbox"/> <i>micafungin</i>	<p>Therapeutic alternatives:</p> <ul style="list-style-type: none"> <li>- <i>anidulafungin</i></li> <li>- <i>caspofungin</i></li> </ul> <p>Powder for injection: 50 mg (as sodium); 100 mg (as sodium) in vial.</p>
<i>potassium iodide</i>	<i>Saturated solution.</i>

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>6.4 Antiviral medicines</b>	
<b>6.4.1 Ant herpes medicines</b>	
<input checked="" type="checkbox"/> aciclovir Therapeutic alternatives: - valaciclovir (oral)	Oral liquid: 200 mg/5 mL [c]. Powder for injection: 250 mg (as sodium salt) in vial. Tablet: 200 mg.
<b>6.4.2 Antiretrovirals</b>	
<p>Based on current evidence and experience of use, medicines in the following classes of antiretrovirals are included as essential medicines for treatment and prevention of HIV (prevention of mother-to-child transmission, pre-exposure prophylaxis (where indicated) and post-exposure prophylaxis). WHO emphasizes the importance of using these products in accordance with global and national guidelines. WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.</p> <p>Scored tablets can be used in children and therefore can be considered for inclusion in the listing of tablets, provided that adequate quality products are available.</p>	
<b>6.4.2.1 Nucleoside/Nucleotide reverse transcriptase inhibitors</b>	
abacavir	Tablet: 300 mg (as sulfate).
lamivudine	Oral liquid: 50 mg/5 mL [c]. Tablet: 150 mg.
tenofovir disoproxil fumarate†	Tablet: 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). † also indicated for pre-exposure prophylaxis.
zidovudine	Capsule: 250 mg. Oral liquid: 50 mg/5 mL. Solution for IV infusion: 10 mg/mL in 20 mL vial. Tablet: 300 mg.
<b>6.4.2.2 Non-nucleoside reverse transcriptase inhibitors</b>	
efavirenz	Tablet: 600 mg.
nevirapine <input checked="" type="checkbox"/>	Oral liquid: 50 mg/5 mL. Tablet: 50 mg (dispersible); 200 mg. <input checked="" type="checkbox"/> > 6 weeks
<b>6.4.2.3 Protease inhibitors</b>	
<p>Selection of protease inhibitor(s) from the Model List will need to be determined by each country after consideration of international and national treatment guidelines and experience. Ritonavir is recommended for use in combination as a pharmacological booster, and not as an antiretroviral in its own right. All other protease inhibitors should be used in boosted forms (e.g. with ritonavir).</p>	
atazanavir + ritonavir	Tablet (heat stable): 300 mg (as sulfate) + 100 mg.
darunavir <input checked="" type="checkbox"/>	Tablet: 75 mg; 400 mg; 600 mg; 800 mg <input checked="" type="checkbox"/> > 3 years
lopinavir + ritonavir	Solid oral dosage form: 40 mg + 10 mg [c]. Tablet (heat stable): 100 mg + 25 mg; 200 mg + 50 mg.
ritonavir	Tablet (heat stable): 25 mg; 100 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>6.4.2.4 Integrase Inhibitors</b>	
dolutegravir <input checked="" type="checkbox"/>	<p>Tablet (dispersible, scored): 10 mg [c].</p> <p><input checked="" type="checkbox"/> ≥ 4 weeks and ≥ 3 kg Tablet: 50 mg</p> <p><input checked="" type="checkbox"/> ≥ 25 kg</p>
raltegravir*	<p>Granules for oral suspension: 100 mg in sachet.</p> <p>Tablet (chewable): 25 mg.</p> <p>Tablet: 400 mg.</p> <p>*For use in pregnant women and in second-line regimens in accordance with WHO treatment guidelines.</p>
<b>6.4.2.5 Fixed-dose combinations of antiretroviral medicines</b>	
abacavir + lamivudine	Tablet (dispersible, scored): 120 mg (as sulfate) + 60 mg.
dolutegravir + lamivudine + tenofovir	Tablet: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
efavirenz + <input checked="" type="checkbox"/> emtricitabine + tenofovir Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
efavirenz + lamivudine + tenofovir	Tablet: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
<input checked="" type="checkbox"/> emtricitabine + tenofovir† Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). † combination also indicated for pre-exposure prophylaxis
lamivudine + zidovudine	Tablet: 30 mg + 60 mg [c]; 150 mg + 300 mg.
<b>6.4.2.6 Medicines for prevention of HIV-related opportunistic infections</b>	
isoniazid + pyridoxine + sulfamethoxazole + trimethoprim	Tablet (scored): 300 mg + 25 mg + 800 mg + 160 mg
<b>6.4.3 Other antiretrovirals</b>	
ribavirin*	<p>Injection for intravenous administration: 800 mg and 1 g in 10 mL phosphate buffer solution.</p> <p>Solid oral dosage form: 200 mg; 400 mg; 600 mg.</p> <p>*For the treatment of viral haemorrhagic fevers</p>
valganciclovir*	<p>Tablet: 450 mg.</p> <p>*For the treatment of cytomegalovirus retinitis (CMVr).</p>

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>Complementary list</b>	
oseltamivir*	Capsule: 30 mg; 45 mg; 75 mg (as phosphate). *Severe illness due to confirmed or suspected influenza virus infection in critically ill hospitalized patients
valganciclovir*[c]	Powder for oral solution: 50 mg/mL Tablet: 450 mg. *For the treatment of cytomegalovirus retinitis (CMV).
<b>6.4.4 Antihepatitis medicines</b>	
<b>6.4.4.1 Medicines for hepatitis B</b>	
<b>6.4.4.1.1 Nucleoside/Nucleotide reverse transcriptase inhibitors</b>	
entecavir	Oral liquid: 0.05 mg/mL Tablet: 0.5 mg; 1 mg
tenofovir disoproxil fumarate	Tablet: 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
<b>6.4.4.2 Medicines for hepatitis C</b>	
Pangenotypic direct-acting antivirals should be considered as therapeutic alternatives for the purposes of selection and procurement at national level.	
<b>6.4.4.2.1 □ Pangenotypic direct-acting antiviral combinations</b>	
daclatasvir*	Tablet: 30 mg; 60 mg (as hydrochloride). *Pangenotypic when used in combination with sofosbuvir
daclatasvir + sofosbuvir	Tablet: 60 mg + 400 mg.
glecaprevir + pibrentasvir	Tablet: 100 mg + 40 mg. Granules: 50 mg + 20 mg in sachet [c].
ravidasvir*	Tablet: 200 mg. *Pangenotypic when used in combination with sofosbuvir
sofosbuvir*	Tablet: 200 mg; 400 mg. *Pangenotypic when used in combination with daclatasvir or ravidasvir
sofosbuvir + velpatasvir	Tablet: 200 mg + 50 mg [c]; 400 mg + 100 mg.
<b>6.4.4.2.2 Non-pangenotypic direct-acting antiviral combinations</b>	
ledipasvir + sofosbuvir	Tablet: 90 mg + 400 mg.
<b>6.4.4.2.3 Other antivirals for hepatitis C</b>	
ribavirin*	Injection for intravenous administration: 800 mg and 1 g in 10 mL phosphate buffer solution. Solid oral dosage form: 200 mg; 400 mg; 600 mg. *For the treatment of hepatitis C, in combination with direct acting anti-viral medicines

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>6.5 Antiprotozoal medicines</b>	
<b>6.5.1 Antamoebic and antigiardiasis medicines</b>	
diloxanide <small>[a]</small>	<b>Tablet:</b> 500 mg (furoate). <small>[a] &gt; 25 kg.</small>
<input type="checkbox"/> metronidazole Therapeutic alternatives: - tinidazole	<b>Injection:</b> 500 mg in 100 mL vial. <b>Oral liquid:</b> 200 mg/5 mL (as benzoate). <b>Tablet:</b> 200 mg; 250 mg; 400 mg; 500 mg.
<b>6.5.2 Antileishmaniasis medicines</b>	
amphotericin B*	<b>Powder for injection:</b> 50 mg (liposomal complex) in vial. <b>Powder for injection:</b> 50 mg (as sodium deoxycholate) in vial. *Liposomal amphotericin B has a better safety profile than the sodium deoxycholate formulation and should be prioritized for selection and use depending on local availability and cost.
meglumine antimoniate	<b>Injection:</b> 1.5 g/5 mL in 5 mL ampoule.
miltefosine	<b>Solid oral dosage form:</b> 10 mg; 50 mg.
paromomycin	<b>Solution for intramuscular injection:</b> 750 mg of paromomycin base (as sulfate).
sodium stibogluconate	<b>Injection:</b> 100 mg/mL in 30 mL vial.
<b>6.5.3 Antimalarial medicines</b>	
<b>6.5.3.1 For curative treatment</b>	
Medicines for the treatment of <i>P. falciparum</i> malaria cases should be used in combination. The list currently recommends combinations according to treatment guidelines. WHO recognizes that not all of the fixed dose combinations (FDCs) in the WHO treatment guidelines exist, and encourages their development and rigorous testing. WHO also encourages development and testing of rectal dosage formulations.	
amodiaquine*	<b>Tablet:</b> 153 mg or 200 mg (as hydrochloride). *To be used in combination with artesunate 50 mg.
artemether*	<b>Oily injection:</b> 80 mg/mL in 1 mL ampoule. *For use in the management of severe malaria.
artemether + lumefantrine*	<b>Tablet:</b> 20 mg + 120 mg. <b>Tablet (dispersible):</b> 20 mg + 120 mg [c]. *Not recommended in the first trimester of pregnancy or in children below 5 kg.

## WHO Model List of Essential Medicines – 23rd List (2023)

artesunate*	<b>Injection:</b> ampoules, containing 60 mg anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution. For use in the management of severe malaria.  <b>Rectal dosage form:</b> 50 mg [c]; 100 mg [c]; 200 mg capsules (for pre-referral treatment of severe malaria only; patients should be taken to an appropriate health facility for follow-up care) [c].  <b>Tablet:</b> 50 mg.  *To be used in combination with either amodiaquine, mefloquine or sulfadoxine + pyrimethamine.
artesunate + amodiaquine*	<b>Tablet:</b> 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg.  *Other combinations that deliver the target doses required such as 153 mg or 200 mg (as hydrochloride) with 50 mg artesunate can be alternatives.
artesunate + mefloquine	<b>Tablet:</b> 25 mg + 55 mg; 100 mg + 220 mg.
artesunate + pyronaridine tetraphosphate □	<b>Granules:</b> 20 mg + 60 mg [c].  <b>Tablet:</b> 60 mg + 180 mg.  □ > 5 kg
chloroquine*	<b>Oral liquid:</b> 50 mg/5 mL (as phosphate or sulfate).  <b>Tablet:</b> 100 mg; 150 mg (as phosphate or sulfate).  *For use only for the treatment of <i>Plasmodium vivax</i> infection.
dihydroartemisinin + piperaquine phosphate □	<b>Tablet:</b> 20 mg + 160 mg; 40 mg + 320 mg.  □ > 5 kg
doxycycline*	<b>Capsule:</b> 100 mg (as hydrochloride or hydiate).  <b>Tablet (dispersible):</b> 100 mg (as monohydrate).  *For use only in combination with quinine.
mefloquine*	<b>Tablet:</b> 250 mg (as hydrochloride).  *To be used in combination with artesunate 50 mg.
primaquine*	<b>Tablet:</b> 7.5 mg; 15 mg (as diphosphate).  *Only for use to achieve radical cure of <i>Plasmodium vivax</i> and <i>Plasmodium ovale</i> infections, given for 14 days.
quinine*	<b>Injection:</b> 300 mg/mL (hydrochloride) in 2 mL ampoule.  <b>Tablet:</b> 300 mg (sulfate) or 300 mg (bisulfate).  *For use only in the management of severe malaria and should be used in combination with doxycycline.
sulfadoxine + pyrimethamine*	<b>Tablet:</b> 500 mg + 25 mg.  *Only in combination with artesunate 50 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

### 6.5.3.2 For chemoprevention

amodiaquine – sulfadoxine + pyrimethamine [c]	<b>Co-packaged dispersible tablets:</b> amodiaquine 76.5 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 250 mg + 12.5 mg [1]; amodiaquine 153 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 500 mg + 25 mg [1].
chloroquine*	<b>Oral liquid:</b> 50 mg/5 mL (as phosphate or sulfate). <b>Tablet:</b> 150 mg (as phosphate or sulfate). *For use only in central American regions, for <i>Plasmodium vivax</i> infections.
doxycycline [a]	<b>Solid oral dosage form:</b> 100 mg (as hydrochloride or hydrate). [a] > 8 years.
mefloquine [a]	<b>Tablet:</b> 250 mg (as hydrochloride). [a] > 5 kg or > 3 months.
proguanil*	<b>Tablet:</b> 100 mg (as hydrochloride). *For use only in combination with chloroquine.
sulfadoxine + pyrimethamine	<b>Tablet:</b> 250 mg + 12.5 mg [c]; 500 mg + 25 mg.

### 6.5.4 AntipROTOZOAL and antitOXOPLASMOSIS medicines

pyrimethamine	<b>Tablet:</b> 25 mg.
sulfadiazine	<b>Tablet:</b> 500 mg.
sulfamethoxazole + trimethoprim	<b>Injection:</b> 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule. <b>Oral liquid:</b> 200 mg + 40 mg/5 mL [c]. <b>Tablet:</b> 100 mg + 20 mg; 400 mg + 80 mg [c]; 800 mg + 160 mg. <b>Tablet (dispersible):</b> 100 mg + 20 mg [c].

#### Complementary List

pentamidine	<b>Tablet:</b> 200 mg; 300 mg (as isethionate).
-------------	---

### 6.5.5 AntitRYPANOSOMAL medicines

fexinidazole*	<b>Tablet:</b> 600 mg *For the treatment of 1 <sup>st</sup> and 2 <sup>nd</sup> stage of human African trypanosomiasis due to <i>Trypanosoma brucei gambiense</i> infection.
<b>Medicines for the treatment of 1<sup>st</sup> stage African trypanosomiasis</b>	
pentamidine*	<b>Powder for injection:</b> 300 mg (as isetionate) in vial. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
suramin sodium*	<b>Powder for injection:</b> 1 g in vial. *To be used for the treatment of the initial phase of <i>Trypanosoma brucei rhodesiense</i> infection.

**WHO Model List of Essential Medicines – 23rd List (2023)**

Medicines for the treatment of 2 <sup>nd</sup> stage African trypanosomiasis	
eflornithine*	Injection: 200 mg/mL (hydrochloride) in 50 mL bottle. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
melarsoprol	Injection: 180 mg/5 mL in 5 mL ampoule (3.6% solution).
nifurtimox *	Tablet (scored): 30 mg; 120 mg. *Only to be used in combination with eflornithine, for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
Complementary List	
melarsoprol [c]	Injection: 180 mg/5 mL in 5 mL ampoule (3.6% solution).
6.5.5.2 American trypanosomiasis	
benznidazole	Tablet: 12.5 mg [c] Tablet (scored): 50 mg; 100 mg.
nifurtimox	Tablet (scored): 30 mg; 120 mg.
6.6 Medicines for ectoparasitic infections	
ivermectin	Tablet: 3 mg
6.7 Medicines for Ebola virus disease	
ansuvimab	Powder for injection: 400 mg
atoltivimab + mabtivimab + odesivimab	Injection: 241.7 mg + 241.7 mg + 241.7 mg in 14.5 mL vial
6.8 Medicines for COVID-19	
WHO recommends that effective and safe therapeutics for prevention and treatment of COVID-19 should be considered as essential medicines in the context of the public health emergency. WHO recommendations are revised and updated regularly in WHO living guidelines for therapeutics for the treatment and prevention of COVID-19.	
Selection of essential therapeutics for COVID-19 at the national level should be informed by recommendations in these guidelines, and consideration of the latest evidence, epidemiology and national priorities.	
The latest WHO Therapeutics and COVID-19: living guideline is available online at: <a href="https://app.magicapp.org/#/guideline/nBkO1E">https://app.magicapp.org/#/guideline/nBkO1E</a>	
The latest WHO Drugs to prevent COVID-19: living guideline is available online at: <a href="https://app.magicapp.org/#/guideline/L6RxYL">https://app.magicapp.org/#/guideline/L6RxYL</a>	

**WHO Model List of Essential Medicines – 23rd List (2023)**
**7. ANTIMIGRAINE MEDICINES**
**7.1 For treatment of acute attack**

acetylsalicylic acid	<b>Tablet:</b> 300 mg to 500 mg.
ibuprofen [c]	<b>Oral liquid:</b> 100 mg/5 mL [c]. <b>Tablet:</b> 200 mg; 400 mg.
paracetamol (acetaminophen)	<b>Oral liquid:</b> 120 mg/5 mL or 125 mg/5 mL*; 250 mg/5 mL [c]. *The presence of both 120 mg/5 mL and 125 mg/5mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided. <b>Suppository:</b> 250 mg [c]. <b>Tablet:</b> 250 mg; 325 mg; 500 mg. <b>Tablet (dispersible):</b> 100 mg, 250 mg [c].
sumatriptan	<b>Tablet:</b> 50 mg

**7.2 For prophylaxis**

<input type="checkbox"/> propranolol Therapeutic alternatives to be reviewed	<b>Tablet:</b> 20 mg; 40 mg (hydrochloride).
---	--

**8. IMMUNOMODULATORS AND ANTI NEOPLASTICS**
**8.1 Immunomodulators for non-malignant disease**
**Complementary List**

<input type="checkbox"/> adalimumab*  Therapeutic alternatives*: - certolizumab pegol - etanercept - golimumab - infliximab  *including quality-assured biosimilars	<b>Injection:</b> 10 mg/0.2 mL [c]; 20 mg/0.4 mL [c]; 40 mg/0.8 mL; 40 mg/0.4 mL.
azathioprine	<b>Oral liquid:</b> 10 mg/mL [c]. <b>Powder for injection:</b> 50 mg [c]; 100 mg (as sodium salt) in vial. <b>Tablet:</b> 25 mg [c]. <b>Tablet (scored):</b> 50 mg.
ciclosporin	<b>Capsule:</b> 25 mg. <b>Concentrate for injection:</b> 50 mg/mL in 1 mL ampoule. <b>Oral liquid:</b> 100 mg/mL [c].
tacrolimus	<b>Capsule (immediate-release):</b> 0.5 mg; 0.75 mg; 1 mg; 2 mg; 5 mg. <b>Granules for oral suspension:</b> 0.2 mg; 1 mg. <b>Injection:</b> 5 mg/mL in 1 mL vial.

## WHO Model List of Essential Medicines – 23rd List (2023)

### 8.2 Antineoplastics and supportive medicines

Medicines listed below should be used according to protocols for treatment of the diseases.

#### 8.2.1 Cytotoxic medicines

##### Complementary List

<i>arsenic trioxide</i>	<b>Concentrate for solution for infusion:</b> 1 mg/mL; 2 mg/mL. – Acute promyelocytic leukaemia
<i>asparaginase*</i> <small>*including quality-assured biosimilars</small>	<b>Powder for injection:</b> 10 000 IU in vial. – Acute lymphoblastic leukaemia.
<i>bendamustine</i>	<b>Injection:</b> 45 mg/0.5 mL; 180 mg/2 mL. – Chronic lymphocytic leukaemia – Follicular lymphoma
<i>bleomycin</i>	<b>Powder for injection:</b> 15 000 IU (as sulfate) in vial. – Hodgkin lymphoma – Kaposi sarcoma – Ovarian germ cell tumour – Testicular germ cell tumour
<i>calcium folinate (leucovorin calcium)</i>	<b>Injection:</b> 3 mg/mL in 10 mL ampoule; 7.5 mg/mL in 2 mL ampoule; 10 mg/mL in 5 mL ampoule. <b>Tablet:</b> 5 mg; 15 mg; 25 mg. – Burkitt lymphoma – Early stage colon cancer – Early stage rectal cancer – Gestational trophoblastic neoplasia – Metastatic colorectal cancer – Osteosarcoma
<i>capecitabine</i>	<b>Tablet:</b> 150 mg; 500 mg. – Early stage colon cancer – Early stage rectal cancer – Metastatic breast cancer – Metastatic colorectal cancer
<i>carboplatin</i>	<b>Injection:</b> 50 mg/5 mL; 150 mg/15 mL; 450 mg/45 mL; 600 mg/60 mL. – Cervical cancer – Early stage breast cancer – Epithelial ovarian cancer – Head and neck cancer (as a radio-sensitizer) – Low-grade glioma – Nasopharyngeal cancer – Nephroblastoma (Wilms tumour) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Retinoblastoma – Testicular germ cell tumour
<i>chlorambucil</i>	<b>Tablet:</b> 2 mg. – Chronic lymphocytic leukaemia

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>cisplatin</i>	<p><b>Injection:</b> 10 mg/10 mL; 20 mg/20 mL; 50 mg/50 mL; 100 mg/100 mL.</p> <ul style="list-style-type: none"> <li>– Cervical cancer</li> <li>– Head and neck cancer (as a radio-sensitizer)</li> <li>– Low-grade glioma</li> <li>– Nasopharyngeal cancer (as a radio-sensitizer)</li> <li>– Non-small cell lung cancer</li> <li>– Osteosarcoma</li> <li>– Ovarian germ cell tumour</li> <li>– Testicular germ cell tumour</li> </ul>
<i>cyclophosphamide</i>	<p><b>Powder for Injection:</b> 500 mg; 1 g; 2 g in vial. <b>Solid oral dosage form:</b> 25 mg; 50 mg.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Chronic lymphocytic leukaemia</li> <li>– Diffuse large B-cell lymphoma</li> <li>– Early stage breast cancer</li> <li>– Ewing sarcoma</li> <li>– Follicular lymphoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Hodgkin lymphoma</li> <li>– Low-grade glioma</li> <li>– Metastatic breast cancer</li> <li>– Multiple myeloma</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>cytarabine</i>	<p><b>Injection:</b> 100 mg/mL in vial <b>Powder for Injection:</b> 100 mg in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Acute myeloid leukaemia</li> <li>– Acute promyelocytic leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Langerhans cell histiocytosis</li> </ul>
<i>dacarbazine</i>	<p><b>Powder for Injection:</b> 100 mg; 200 mg in vial.</p> <ul style="list-style-type: none"> <li>– Hodgkin lymphoma</li> </ul>
<i>dactinomycin</i>	<p><b>Powder for Injection:</b> 500 micrograms in vial.</p> <ul style="list-style-type: none"> <li>– Ewing sarcoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>daunorubicin</i>	<p><b>Injection:</b> 2 mg/mL; 5 mg/mL (as hydrochloride) in vial. <b>Powder for Injection:</b> 20 mg; 50 mg (as hydrochloride) in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Acute myeloid leukaemia</li> <li>– Acute promyelocytic leukaemia</li> </ul>

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>docetaxel</i>	<p><b>Injection:</b> 20 mg/mL; 40 mg/mL.</p> <ul style="list-style-type: none"> <li>– Early stage breast cancer</li> <li>– Metastatic breast cancer</li> <li>– Metastatic prostate cancer</li> </ul>
<i>doxorubicin</i>	<p><b>Injection:</b> 2 mg/mL (hydrochloride) in vial.</p> <p><b>Powder for Injection:</b> 10 mg; 50 mg (hydrochloride) in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Diffuse large B-cell lymphoma</li> <li>– Early stage breast cancer</li> <li>– Ewing sarcoma</li> <li>– Follicular lymphoma</li> <li>– Hodgkin lymphoma</li> <li>– Kaposi sarcoma</li> <li>– Metastatic breast cancer</li> <li>– Multiple myeloma</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Osteosarcoma</li> </ul>
<i>doxorubicin (as pegylated liposomal)</i>	<p><b>Injection:</b> 2 mg/mL (hydrochloride) in 10 mL, 25 mL vial</p> <ul style="list-style-type: none"> <li>– Kaposi sarcoma</li> </ul>
<i>etoposide</i>	<p><b>Capsule:</b> 50 mg, 100 mg.</p> <p><b>Injection:</b> 20 mg/mL in 5 mL ampoule.</p> <p><b>Powder for Injection:</b> 100 mg (as phosphate) in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Acute myeloid leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Ewing sarcoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Hodgkin lymphoma</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Non-small cell lung cancer</li> <li>– Osteosarcoma</li> <li>– Ovarian germ cell tumour</li> <li>– Retinoblastoma</li> <li>– Testicular germ cell tumour</li> </ul>
<i>fudarabine</i>	<p><b>Powder for Injection:</b> 50 mg (phosphate) in vial.</p> <p><b>Tablet:</b> 10 mg</p> <ul style="list-style-type: none"> <li>– Chronic lymphocytic leukaemia.</li> </ul>
<i>fluorouracil</i>	<p><b>Injection:</b> 50 mg/mL in vial.</p> <ul style="list-style-type: none"> <li>– Early stage breast cancer</li> <li>– Early stage colon cancer</li> <li>– Early stage rectal cancer</li> <li>– Metastatic colorectal cancer</li> <li>– Nasopharyngeal cancer</li> </ul>
<i>gemcitabine</i>	<p><b>Powder for Injection:</b> 200 mg; 1 g in vial.</p> <ul style="list-style-type: none"> <li>– Epithelial ovarian cancer</li> <li>– Non-small cell lung cancer</li> </ul>

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>hydroxycarbamide (hydroxyurea)</i>	<b>Solid oral dosage form:</b> 100 mg [c]; 200 mg; 300 mg; 400 mg; 500 mg; 1 g. – Chronic myeloid leukaemia
<i>ifosfamide</i>	<b>Powder for injection:</b> 500 mg; 1 g; 2 g in vial. – Anaplastic large cell lymphoma – Burkitt lymphoma – Ewing sarcoma – Nephroblastoma (Wilms tumour) – Ovarian germ cell tumour – Osteosarcoma – Rhabdomyosarcoma – Testicular germ cell tumour
<i>irinotecan</i>	<b>Injection:</b> 40 mg/2 mL in 2 mL vial; 100 mg/5 mL in 5 mL vial; 500 mg/25 mL in 25 mL vial. – Metastatic colorectal cancer – Nephroblastoma (Wilms tumour) – Rhabdomyosarcoma
<i>melphalan</i>	<b>Tablet:</b> 2 mg <b>Powder for injection:</b> 50 mg in vial – Multiple myeloma
<i>mercaptopurine</i>	<b>Tablet:</b> 50 mg. <b>Oral liquid:</b> 20 mg/mL [c]. – Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia – Langerhans cell histiocytosis
<i>methotrexate</i>	<b>Concentrated injection:</b> 1000 mg/10 mL. <b>Injection:</b> 50mg/2 mL. <b>Powder for injection:</b> 50 mg (as sodium) in vial. <b>Tablet:</b> 2.5 mg (as sodium). – Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Early stage breast cancer – Gestational trophoblastic neoplasia – Langerhans cell histiocytosis – Osteosarcoma
<i>oxaliplatin</i>	<b>Injection:</b> 50 mg/10 mL in 10 mL vial; 100 mg/20 mL in 20 mL vial; 200 mg/40 mL in 40 mL vial. <b>Powder for injection:</b> 50 mg; 100 mg in vial. – Early stage colon cancer – Metastatic colorectal cancer

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>paclitaxel</i>	<b>Injection:</b> 6 mg/mL in vial. <ul style="list-style-type: none"> <li>– Cervical cancer</li> <li>– Epithelial ovarian cancer</li> <li>– Early stage breast cancer</li> <li>– Metastatic breast cancer</li> <li>– Kaposi sarcoma</li> <li>– Nasopharyngeal cancer</li> <li>– Non-small cell lung cancer</li> <li>– Ovarian germ cell tumour</li> </ul>
<i>pegaspargase*</i> <small>*Including quality-assured biosimilars</small>	<b>Injection:</b> 3750 units/5 mL in vial. <b>Powder for injection:</b> 3750 units in vial. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> </ul>
<i>procarbazine [c]</i>	<b>Capsule:</b> 50 mg (as hydrochloride). <ul style="list-style-type: none"> <li>– Hodgkin lymphoma</li> </ul>
<i>realgar-Indigo naturalis formulation</i>	<b>Tablet:</b> 270 mg (containing tetra-arsenic tetra-sulfide 30 mg). <ul style="list-style-type: none"> <li>– Acute promyelocytic leukaemia</li> </ul>
<i>tioguanine [c]</i>	<b>Solid oral dosage form:</b> 40 mg. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> </ul>
<i>vinblastine</i>	<b>Injection:</b> 10 mg/10 mL (sulfate) in vial. <b>Powder for injection:</b> 10 mg (sulfate) in vial. <ul style="list-style-type: none"> <li>– Anaplastic large cell lymphoma</li> <li>– Hodgkin lymphoma</li> <li>– Kaposi sarcoma</li> <li>– Langerhans cell histiocytosis</li> <li>– Low-grade glioma</li> <li>– Ovarian germ cell tumour</li> <li>– Testicular germ cell tumour</li> </ul>
<i>vincristine</i>	<b>Injection:</b> 1 mg/mL (sulfate); 2 mg/2 mL (sulfate) in vial. <b>Powder for injection:</b> 1 mg; 5 mg (sulfate) in vial. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Burkitt lymphoma</li> <li>– Diffuse large B-cell lymphoma</li> <li>– Ewing sarcoma</li> <li>– Follicular lymphoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Hodgkin lymphoma</li> <li>– Kaposi sarcoma</li> <li>– Langerhans cell histiocytosis</li> <li>– Low-grade glioma</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Retinoblastoma</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>vinorelbine</i>	<b>Capsule:</b> 20 mg; 30 mg; 80 mg. <b>Injection:</b> 10 mg/mL in 1 mL, 5 mL vial. <ul style="list-style-type: none"> <li>– Non-small cell lung cancer</li> <li>– Metastatic breast cancer</li> <li>– Rhabdomyosarcoma</li> </ul>

**WHO Model List of Essential Medicines – 23rd List (2023)**

8.2.2 Targeted therapies	
Complementary List	
<i>all-trans-retinoid acid (ATRA)</i>	<b>Capsule:</b> 10 mg. – Acute promyelocytic leukaemia.
<i>bortezomib</i>	<b>Powder for injection:</b> 3.5 mg in vial. – Multiple myeloma
<i>dasatinib</i>	<b>Tablet:</b> 20 mg; 50 mg; 70 mg; 80 mg; 100 mg; 140 mg. – Imatinib-resistant chronic myeloid leukaemia
<input checked="" type="checkbox"/> <i>erlotinib</i> <i>Therapeutic alternatives:</i> – afatinib – gefitinib	<b>Tablet:</b> 100 mg, 150 mg. – EGFR mutation-positive advanced non-small cell lung cancer
<i>everolimus</i>	<b>Tablet:</b> 2.5 mg; 5 mg; 7.5 mg; 10 mg. <b>Tablet (dispersible):</b> 2 mg; 3 mg; 5 mg. – Subependymal giant cell astrocytoma
<i>ibrutinib</i>	<b>Capsule:</b> 140 mg. – Relapsed/refractory chronic lymphocytic leukaemia
<i>imatinib</i>	<b>Solid oral dosage form:</b> 100 mg; 400 mg. – Chronic myeloid leukaemia – Gastrointestinal stromal tumour – Philadelphia chromosome positive acute lymphoblastic leukaemia
<i>nilotinib</i>	<b>Capsule:</b> 150 mg; 200 mg. – Imatinib-resistant chronic myeloid leukaemia
<i>rituximab*</i> *including quality-assured biosimilars	<b>Injection (intravenous):</b> 100 mg/10 mL in 10 mL vial; 500 mg/50 mL in 50 mL vial. – Burkitt lymphoma – Diffuse large B-cell lymphoma – Chronic lymphocytic leukaemia – Follicular lymphoma
<i>trastuzumab*</i> *including quality-assured biosimilars	<b>Powder for injection:</b> 60 mg; 150 mg; 440 mg in vial. – Early stage HER2-positive breast cancer – Metastatic HER2-positive breast cancer

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>8.2.3 Immunomodulators</b>	
<b>Complementary List</b>	
<i>filgrastim*</i> *including quality-assured biosimilars	<b>Injection:</b> 120 micrograms/0.2 mL; 300 micrograms/0.5 mL; 480 micrograms/0.8 mL in pre-filled syringe.  <b>Injection:</b> 300 micrograms/mL in 1 mL vial; 480 micrograms/1.6 mL in 1.6 mL vial.  – Primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy. – Secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy – To facilitate administration of dose dense chemotherapy regimens
<i>lenalidomide</i>	<b>Capsule:</b> 25 mg.  – Multiple myeloma
<input checked="" type="checkbox"/> <i>nivolumab*</i> Therapeutic alternatives*: - pembrolizumab *including quality-assured biosimilars	<b>Concentrate solution for infusion:</b> 10 mg/mL  – Metastatic melanoma
<i>pegfilgrastim*</i> *including quality-assured biosimilars	<b>Injection:</b> 6 mg/0.6 mL in pre-filled syringe.  – Primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy – Secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy – To facilitate administration of dose dense chemotherapy regimens
<i>thalidomide</i>	<b>Capsule:</b> 50 mg.  – Multiple myeloma
<b>8.2.4 Hormones and antihormones</b>	
<b>Complementary List</b>	
<input checked="" type="checkbox"/> <i>abiraterone</i> Therapeutic alternatives: - enzalutamide	<b>Tablet:</b> 250 mg; 500 mg.  – Metastatic castration-resistant prostate cancer
<input checked="" type="checkbox"/> <i>anastrozole</i> Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (L02BG Aromatase Inhibitors)	<b>Tablet:</b> 1 mg.  – Early stage breast cancer – Metastatic breast cancer
<input checked="" type="checkbox"/> <i>bicalutamide</i> Therapeutic alternatives: - flutamide - nilutamide	<b>Tablet:</b> 50 mg.  – Metastatic prostate cancer

## WHO Model List of Essential Medicines – 23rd List (2023)

dexamethasone	<b>Injection:</b> 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule. <b>Oral liquid:</b> 2 mg/5 mL [c]. <b>Tablet:</b> 2 mg [c]; 4 mg. – Acute lymphoblastic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Multiple myeloma
hydrocortisone	<b>Powder for Injection:</b> 100 mg (as sodium succinate) in vial. – Acute lymphoblastic leukaemia – Burkitt lymphoma
<input checked="" type="checkbox"/> leuprorelin  Therapeutic alternatives: - goserelin - triptorelin	<b>Injection:</b> 7.5 mg; 22.5 mg in pre-filled syringe. – Early stage breast cancer – Metastatic prostate cancer.
methylprednisolone [c]	<b>Injection:</b> 40 mg/mL (as sodium succinate) in 1 mL single-dose vial and 5 mL multi-dose vials; 80 mg/mL (as sodium succinate) in 1 mL single-dose vial. – Acute lymphoblastic leukemia – Burkitt lymphoma
<input checked="" type="checkbox"/> prednisolone  Therapeutic alternatives: - prednisone	<b>Oral liquid:</b> 5 mg/mL [c]. <b>Tablet:</b> 5 mg; 25 mg. – Acute lymphoblastic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Chronic lymphocytic leukaemia – Diffuse large B-cell lymphoma – Follicular lymphoma – Hodgkin lymphoma – Langerhans cell histiocytosis – Metastatic castration-resistant prostate cancer – Multiple myeloma
tamoxifen	<b>Tablet:</b> 10 mg; 20 mg (as citrate). – Early stage breast cancer – Metastatic breast cancer.

**8.2.5 Supportive medicines**

<b>Complementary List</b>	
allopurinol [c]	<b>Tablet:</b> 100 mg; 300 mg. – Tumour lysis syndrome
mesna	<b>Injection:</b> 100 mg/mL in 4 mL and 10 mL ampoules. <b>Tablet:</b> 400 mg; 600 mg. – Burkitt lymphoma – Ewing sarcoma – Nephroblastoma (Wilms tumour) – Ovarian germ cell tumour – Osteosarcoma – Rhabdomyosarcoma – Testicular germ cell tumour

## WHO Model List of Essential Medicines – 23rd List (2023)

<i>rasburicase</i>	<i>Powder and solvent for solution for infusion: 1.5 mg; 7.5 mg in vial.</i> – Tumour lysis syndrome
<i>zoledronic acid</i>	<i>Concentrate solution for infusion: 4 mg/5 mL in 5 mL vial.</i> <i>Solution for infusion: 4 mg/100 mL in 100 mL bottle.</i> – Malignancy-related bone disease

## 9. THERAPEUTIC FOODS

ready-to-use therapeutic food [c]	<i>Biscuit or paste*.</i> *of nutritional composition as determined by the UN joint statement on the community-based management of severe acute malnutrition and Codex alimentarius guidelines.
-----------------------------------	--

## 10. MEDICINES AFFECTING THE BLOOD

## 10.1 Ant anaemia medicines

ferrous salt	<i>Oral liquid:</i> equivalent to 25 mg iron (as sulfate)/mL. <i>Tablet:</i> equivalent to 60 mg iron.
ferrous salt + folic acid	<i>Tablet:</i> equivalent to 60 mg elemental iron + 400 micrograms folic acid.* *nutritional supplement for use during pregnancy <i>Tablet:</i> equivalent to 60 mg elemental iron + 2.8 mg folic acid.** **for weekly iron and folic acid supplementation
folic acid	<i>Tablet:</i> 400 micrograms*; 1 mg; 5 mg. *periconceptual use for prevention of first occurrence of neural tube defects
hydroxocobalamin	<i>Injection:</i> 1 mg/mL (as acetate, as hydrochloride or as sulfate) in 1 mL ampoule.

*Complementary List*

<input type="checkbox"/> <i>erythropoiesis-stimulating agents*</i> Therapeutic alternatives: - epoetin alfa, beta and theta - darbepoetin alfa - methoxy polyethylene glycol-epoetin beta *including quality-assured biosimilars	<i>Injection: pre-filled syringe</i> 1000 IU/0.5 mL; 2000 IU/0.5 mL; 3000 IU/0.3 mL; 4000 IU/0.4 mL; 5000 IU/0.5 mL; 6000 IU/0.6 mL; 8000 IU/0.8 mL; 10 000 IU/1 mL; 20 000 IU/0.5 mL; 40 000 IU/1 mL.
<b>10.2 Medicines affecting coagulation</b>	
<input type="checkbox"/> dabigatran Therapeutic alternatives: - apixaban - edoxaban - rivaroxaban	<i>Capsule:</i> 110 mg; 150 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

<input checked="" type="checkbox"/> enoxaparin*	<b>Injection:</b> ampoule or pre-filled syringe 20 mg/0.2 mL; 40 mg/0.4 mL; 60 mg/0.6 mL; 80 mg/0.8 mL; 100 mg/1 mL; 120 mg/0.8 mL; 150 mg/1 mL.
<b>*including quality-assured biosimilars</b>	
<input checked="" type="checkbox"/> heparin sodium	<b>Injection:</b> 1000 IU/mL; 5000 IU/mL; 20 000 IU/mL in 1 mL ampoule.
<input checked="" type="checkbox"/> phytomenadione	<b>Injection:</b> 1 mg/mL [c]; 10 mg/mL in ampoule. <b>Tablet:</b> 10 mg.
<input checked="" type="checkbox"/> protamine sulfate	<b>Injection:</b> 10 mg/mL in 5 mL ampoule.
<input checked="" type="checkbox"/> tranexamic acid	<b>Injection:</b> 100 mg/mL in 10 mL ampoule.
<input checked="" type="checkbox"/> warfarin	<b>Tablet:</b> 1 mg; 2 mg; 5 mg (sodium).
<b>Therapeutic alternatives to be reviewed</b>	
<b>Complementary List</b>	
<input checked="" type="checkbox"/> desmopressin [c]	<b>Injection:</b> 4 micrograms/mL (as acetate) in 1 mL ampoule. <b>Nasal spray:</b> 10 micrograms (as acetate) per dose.
<input checked="" type="checkbox"/> heparin sodium [c]	<b>Injection:</b> 1000 IU/mL; 5000 IU/mL in 1 mL ampoule.
<input checked="" type="checkbox"/> protamine sulfate [c]	<b>Injection:</b> 10 mg/mL in 5 mL ampoule.
<input checked="" type="checkbox"/> warfarin [c]	<b>Tablet:</b> 0.5 mg; 1 mg; 2 mg; 5 mg (sodium).
<b>Therapeutic alternatives to be reviewed</b>	
<b>10.3 Other medicines for haemoglobinopathies</b>	
<input checked="" type="checkbox"/> deferasirox	<b>Tablet (dispersible):</b> 100 mg; 125 mg; 250 mg; 400 mg; 500 mg. <b>Tablet (film-coated):</b> 90 mg; 180 mg; 360 mg.
<b>Complementary List</b>	
<input checked="" type="checkbox"/> deferoxamine	<b>Powder for injection:</b> 500 mg (mesilate) in vial.
<input checked="" type="checkbox"/> hydroxycarbamide (hydroxyurea)	<b>Solid oral dosage form:</b> 100 mg [c]; 200 mg; 500 mg; 1 g.
<b>11. BLOOD PRODUCTS OF HUMAN ORIGIN AND PLASMA SUBSTITUTES</b>	
<b>11.1 Blood and blood components</b>	
In accordance with the World Health Assembly resolution WHA63.12, WHO recognizes that achieving self-sufficiency, unless special circumstances preclude it, in the supply of safe blood components based on voluntary, non-remunerated blood donation, and the security of that supply are important national goals to prevent blood shortages and meet the transfusion requirements of the patient population. All preparations should comply with the WHO requirements.	
<input checked="" type="checkbox"/> cryoprecipitate, pathogen-reduced	<b>Injection:</b> frozen liquid in bag or lyophilized powder in vial containing: - > 50 IU Factor VIII - > 100 IU vWF - > 140 mg clottable fibrinogen per unit
<b>Therapeutic alternatives:</b>	
- cryoprecipitate (not pathogen-reduced)	
<input checked="" type="checkbox"/> fresh-frozen plasma	
<input checked="" type="checkbox"/> platelets	
<input checked="" type="checkbox"/> red blood cells	

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>whole blood</b>	
<b>11.2 Plasma-derived medicines</b>	
All human plasma-derived medicines should comply with the WHO requirements.	
<b>11.2.1 Human immunoglobulins</b>	
anti-D immunoglobulin	Injection: 250 micrograms in single-dose vial.
anti-rabies Immunoglobulin	Injection: 150 IU/mL in vial.
anti-tetanus Immunoglobulin	Injection: 500 IU in vial.
<b>Complementary List</b>	
normal immunoglobulin	<p><i>Intramuscular administration:</i> 16% protein solution.</p> <p><i>Subcutaneous administration:</i> 15%; 16% protein solution.</p> <ul style="list-style-type: none"> <li>- Primary immune deficiency.</li> </ul> <p><i>Intravenous administration:</i> 5%; 10% protein solution.</p> <ul style="list-style-type: none"> <li>- Primary immune deficiency</li> <li>- Kawasaki disease</li> <li>- Langerhans cell histiocytosis</li> </ul>
<b>11.2.2 Blood coagulation factors</b>	
<b>Complementary List</b>	
coagulation factor VIII	Powder for injection: 250 IU; 500 IU; 1000 IU in vial.
<input type="checkbox"/> coagulation factor IX	
Therapeutic alternatives: - coagulation factor IX complex	Powder for injection: 500 IU; 1000 IU in vial.
<b>11.3 Plasma substitutes</b>	
<input type="checkbox"/> dextran 70	
Therapeutic alternatives: - polygeline injectable solution 3.5%	Injectable solution: 6%.
<b>12. CARDIOVASCULAR MEDICINES</b>	
<b>12.1 Antianginal medicines</b>	
<input type="checkbox"/> bisoprolol	
Therapeutic alternatives: - carvedilol - metoprolol	Tablet: 1.25 mg; 5 mg.
glyceryl trinitrate	Tablet (sublingual): 500 micrograms.
Isosorbide dinitrate	Tablet (sublingual): 5 mg.
verapamil	Tablet: 40 mg; 80 mg (hydrochloride).

## WHO Model List of Essential Medicines – 23rd List (2023)

12.2 Antiarrhythmic medicines	
<input checked="" type="checkbox"/> bisoprolol	Therapeutic alternatives: - carvedilol - metoprolol
	Tablet: 1.25 mg; 5 mg.
digoxin	Injection: 250 micrograms/mL in 2 mL ampoule. Oral liquid: 50 micrograms/mL. Tablet: 62.5 micrograms; 250 micrograms.
epinephrine (adrenaline)	Injection: 100 micrograms/mL (as acid tartrate or hydrochloride) in 10 mL ampoule.
lidocaine	Injection: 20 mg/mL (hydrochloride) in 5 mL ampoule.
verapamil	Injection: 2.5 mg/mL (hydrochloride) in 2 mL ampoule. Tablet: 40 mg; 80 mg (hydrochloride).
<i>Complementary List</i>	
amiodarone	Injection: 50 mg/mL (hydrochloride) in 3 mL ampoule. Tablet: 100 mg; 200 mg; 400 mg (hydrochloride).
12.3 Antihypertensive medicines	
<input checked="" type="checkbox"/> amlodipine	Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives)
	Tablet: 5 mg (as maleate, mesylate or besylate).
<input checked="" type="checkbox"/> bisoprolol	Therapeutic alternatives: - atenolol* - carvedilol - metoprolol
	Tablet: 1.25 mg; 5 mg. *atenolol should not be used as a first-line agent in uncomplicated hypertension in patients > 60 years
<input checked="" type="checkbox"/> enalapril	Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain)
	Oral liquid: 1 mg/mL (as hydrogen maleate) [c]. Tablet: 2.5 mg; 5 mg; 10 mg (as hydrogen maleate).
hydralazine*	Powder for Injection: 20 mg (hydrochloride) in ampoule. Tablet: 25 mg; 50 mg (hydrochloride). *Hydralazine is listed for use only in the acute management of severe pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
<input checked="" type="checkbox"/> hydrochlorothiazide	Therapeutic alternatives: - chlorothiazide - chlorthalidone - indapamide
	Oral liquid: 50 mg/5 mL. Solid oral dosage form: 12.5 mg; 25 mg.

## WHO Model List of Essential Medicines – 23rd List (2023)

<input checked="" type="checkbox"/> lisinopril + <input type="checkbox"/> amlodipine	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril)	Tablet: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg.
- 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	
<input type="checkbox"/> lisinopril + <input type="checkbox"/> hydrochlorothiazide	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril)	Tablet: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg.
- chlorthalidone, chlorothiazide, Indapamide (for hydrochlorothiazide)	
<input type="checkbox"/> losartan	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	Tablet: 25 mg; 50 mg; 100 mg.
<input type="checkbox"/> methyldopa*	
methyldopa*	*Methyldopa is listed for use only in the management of pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
<input type="checkbox"/> telmisartan + <input type="checkbox"/> amlodipine	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan)	Tablet: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg.
- 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	
<input type="checkbox"/> telmisartan + <input type="checkbox"/> hydrochlorothiazide	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan)	Tablet: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg.
- chlorthalidone, chlorothiazide, Indapamide (for hydrochlorothiazide)	
<b>Complementary List</b>	
sodium nitroprusside	Powder for Infusion: 50 mg in ampoule.

## WHO Model List of Essential Medicines – 23rd List (2023)

12.4 Medicines used in heart failure	
<input type="checkbox"/> bisoprolol	
Therapeutic alternatives:	
- carvedilol	
- metoprolol	
digoxin	<b>Injection:</b> 250 micrograms/mL in 2 mL ampoule. <b>Oral liquid:</b> 50 micrograms/mL. <b>Tablet:</b> 62.5 micrograms; 250 micrograms.
<input type="checkbox"/> enalapril	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain)	<b>Tablet:</b> 2.5 mg; 5 mg; 10 mg (as hydrogen maleate).
<input type="checkbox"/> furosemide	
Therapeutic alternatives:	
- bumetanide	
- torasemide	<b>Injection:</b> 10 mg/mL in 2 mL, 5 mL ampoule. <b>Oral liquid:</b> 20 mg/5 mL; 50 mg/5 mL [c]. <b>Tablet:</b> 20 mg; 40 mg.
<input type="checkbox"/> hydrochlorothiazide	
Therapeutic alternatives:	
- chlorothiazide	
- chlorthalidone	
- indapamide	<b>Oral liquid:</b> 50 mg/5 mL. <b>Solid oral dosage form:</b> 25 mg.
<input type="checkbox"/> losartan	
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	<b>Tablet:</b> 25 mg; 50 mg; 100 mg.
spironolactone	<b>Tablet:</b> 25 mg.
Complementary List	
<input type="checkbox"/> digoxin [c]	<b>Injection:</b> 100 micrograms/mL in 1 mL ampoule; 250 micrograms/mL in 2 mL ampoule. <b>Oral liquid:</b> 50 micrograms/mL. <b>Tablet:</b> 62.5 micrograms; 125 micrograms; 250 mg micrograms.
dopamine	<b>Injection:</b> 40 mg/mL (hydrochloride) in 5 mL vial.
12.5 Antithrombotic medicines	
12.5.1 Anti-platelet medicines	
acetylsalicylic acid	<b>Tablet:</b> 100 mg.
clopidogrel	<b>Tablet:</b> 75 mg; 300 mg
12.5.2 Thrombolytic medicines	
Complementary List	
alteplase	<b>Powder for injection:</b> 10 mg; 20 mg; 50 mg in vial
streptokinase	<b>Powder for injection:</b> 1.5 million IU in vial.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>12.6 Lipid-lowering agents</b>	
<input type="checkbox"/> simvastatin*	<p>Therapeutic alternatives:</p> <ul style="list-style-type: none"> <li>- atorvastatin</li> <li>- fluvastatin</li> <li>- lovastatin</li> <li>- pravastatin</li> </ul> <p><b>Tablet:</b> 5 mg; 10 mg; 20 mg; 40 mg.</p> <p>*For use in high-risk patients.</p>
<b>12.7 Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease</b>	
<input type="checkbox"/> acetylsalicylic acid + <input type="checkbox"/> atorvastatin + <input type="checkbox"/> ramipril	<p>Therapeutic alternatives:</p> <ul style="list-style-type: none"> <li>- fluvastatin, lovastatin, pravastatin, simvastatin (for atorvastatin)</li> <li>- 4<sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for ramipril)</li> </ul> <p><b>Tablet:</b> 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg.</p>
<input type="checkbox"/> acetylsalicylic acid + <input type="checkbox"/> simvastatin + <input type="checkbox"/> ramipril + <input type="checkbox"/> atenolol + <input type="checkbox"/> hydrochlorothiazide	<p>Therapeutic alternatives:</p> <ul style="list-style-type: none"> <li>- atorvastatin, fluvastatin, lovastatin, pravastatin (for simvastatin)</li> <li>- 4<sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for ramipril)</li> <li>- bisoprolol, carvedilol, metoprolol (for atenolol)</li> <li>- chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)</li> </ul> <p><b>Tablet:</b> 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg.</p>
<input type="checkbox"/> atorvastatin + <input type="checkbox"/> perindopril + <input type="checkbox"/> amlodipine	<p>Therapeutic alternatives:</p> <ul style="list-style-type: none"> <li>- fluvastatin, lovastatin, pravastatin, simvastatin (for atorvastatin)</li> <li>- 4<sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for perindopril)</li> <li>- 4<sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)</li> </ul> <p><b>Tablet:</b> 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg.</p>
<b>13. DERMATOLOGICAL MEDICINES</b>	
<b>13.1 Antifungal medicines</b>	
<input type="checkbox"/> miconazole	Cream or ointment: 2% (nitrate).
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (D01AC Imidazole and triazole derivatives) excluding combinations	
selenium sulfide	Detergent-based suspension: 2%.
sodium thiosulfate	Solution: 15%.
terbinafine	Cream or ointment: 1% (hydrochloride).

**WHO Model List of Essential Medicines – 23rd List (2023)**

13.2 Anti-Infective medicines	
mupirocin	Cream: 2% (as calcium). Ointment: 2%.
potassium permanganate	Aqueous solution: 1:10 000.
silver sulfadiazine <input checked="" type="checkbox"/>	Cream: 1%. <input checked="" type="checkbox"/> > 2 months.
13.3 Anti-Inflammatory and antipruritic medicines	
<input checked="" type="checkbox"/> betamethasone <input checked="" type="checkbox"/>	Cream or ointment: 0.1% (as valerate). <input checked="" type="checkbox"/> Hydrocortisone preferred in neonates.
Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D07AC Corticosteroids, potent (group III))	
calamine	Lotion.
<input checked="" type="checkbox"/> hydrocortisone	Cream or ointment: 1% (acetate).
Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D07AA Corticosteroids, weak (group I))	
13.4 Medicines affecting skin differentiation and proliferation	
benzoyl peroxide	Cream or lotion: 5%.
<input checked="" type="checkbox"/> calcipotriol	Cream or ointment: 50 micrograms/mL (0.005%). Lotion: 50 micrograms/mL (0.005%).
Therapeutic alternatives: - calcitriol - tacalcitol	
coal tar	Solution: 5%.
fluorouracil	Ointment: 5%.
<input checked="" type="checkbox"/> podophyllum resin	Solution: 10% to 25%.
Therapeutic alternatives: - podophyllotoxin	
salicylic acid	Solution: 5%.
urea	Cream or ointment: 5%; 10%.
Complementary List	
methotrexate	Tablet: 2.5 mg; 10 mg (as sodium).
13.5 Scabicides and pediculicides	
<input checked="" type="checkbox"/> benzyl benzoate <input checked="" type="checkbox"/>	Lotion: 25%. <input checked="" type="checkbox"/> > 2 years.
Therapeutic alternatives: - precipitated sulfur topical ointment	
permethrin	Cream: 5%. Lotion: 1%.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>14. DIAGNOSTIC AGENTS</b>	
<b>14.1 Ophthalmic medicines</b>	
fluorescein	<b>Eye drops:</b> 1% (sodium salt).
<input type="checkbox"/> tropicamide Therapeutic alternatives: - atropine - cyclopentolate	<b>Eye drops:</b> 0.5%.
<b>14.2 Radiocontrast media</b>	
<input type="checkbox"/> amidotrizoate Therapeutic alternatives to be reviewed	<b>Injection:</b> 140 mg to 420 mg iodine/mL (as sodium or meglumine salt) in 20 mL ampoule.
barium sulfate	<b>Aqueous suspension.</b>
<input type="checkbox"/> iohexol Therapeutic alternatives to be reviewed	<b>Injection:</b> 140 mg to 350 mg iodine/mL in 5 mL, 10 mL, 20 mL ampoules.
<i>Complementary List</i>	
<input type="checkbox"/> barium sulfate [c]	<b>Aqueous suspension.</b>
<input type="checkbox"/> meglumine iotroxate Therapeutic alternatives to be reviewed	<b>Solution:</b> 5 g to 8 g iodine in 100 mL to 250 mL
<b>15. ANTISEPTICS AND DISINFECTANTS</b>	
<b>15.1 Antiseptics</b>	
<input type="checkbox"/> chlorhexidine Therapeutic alternatives to be reviewed	<b>Solution:</b> 5% (digluconate).
<input type="checkbox"/> ethanol Therapeutic alternatives: - propanol	<b>Solution:</b> 70% (denatured).
<input type="checkbox"/> povidone iodine Therapeutic alternatives: - iodine	<b>Solution:</b> 10% (equivalent to 1% available iodine).
<b>15.2 Disinfectants</b>	
alcohol based hand rub	<b>Solution:</b> containing ethanol 80% volume/volume. <b>Solution:</b> containing isopropyl alcohol 75% volume/volume.
chlorine base compound	<b>Liquid:</b> (0.1% available chlorine) for solution. <b>Powder:</b> (0.1% available chlorine) for solution. <b>Solid:</b> (0.1% available chlorine) for solution.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<input checked="" type="checkbox"/> chloroxylenol	Solution: 4.8%.
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (D08AE Phenol and derivatives)	
<input checked="" type="checkbox"/> glutaral	Solution: 2%.
<b>16. DIURETICS</b>	
<input checked="" type="checkbox"/> amiloride	Tablet: 5 mg (hydrochloride).
<input checked="" type="checkbox"/> furosemide	Injection: 10 mg/mL in 2 mL, 5 mL ampoule. Oral liquid: 20 mg/5 mL; 50 mg/5 mL [c]. Tablet: 20 mg; 40 mg.
Therapeutic alternatives:	
- bumetanide - torasemide	
<input checked="" type="checkbox"/> hydrochlorothiazide	Solid oral dosage form: 25 mg.
Therapeutic alternatives:	
- chlorothiazide - chlortalidone - indapamide	
mannitol	Injectable solution: 10%; 20%.
spironolactone	Tablet: 25 mg.
<i>Complementary List</i>	
<input checked="" type="checkbox"/> hydrochlorothiazide[c]	
Therapeutic alternatives:	
- chlorothiazide - chlortalidone	Tablet (scored): 25 mg.
mannitol [c]	Injectable solution: 10%; 20%.
spironolactone[c]	Oral liquid: 5 mg/5 mL; 10 mg/5 mL; 25 mg/5 mL. Tablet: 25 mg.
<b>17. GASTROINTESTINAL MEDICINES</b>	
<i>Complementary List</i>	
pancreatic enzymes[c]	Age-appropriate formulations and doses including lipase, protease and amylase.
<b>17.1 Antidiarrhoeal medicines</b>	
<input checked="" type="checkbox"/> omeprazole	Powder for injection: 40 mg in vial Powder for oral liquid: 20 mg; 40 mg sachets. Solid oral dosage form: 10 mg; 20 mg; 40 mg.
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (A02BC Proton pump inhibitors) excluding combinations	
<input checked="" type="checkbox"/> ranitidine	Injection: 25 mg/mL (as hydrochloride) in 2 mL ampoule. Oral liquid: 75 mg/5 mL (as hydrochloride). Tablet: 150 mg (as hydrochloride).
Therapeutic alternatives:	
- 4 <sup>th</sup> level ATC chemical subgroup (A02BA H <sub>2</sub> -receptor antagonists) excluding combinations	

## WHO Model List of Essential Medicines – 23rd List (2023)

### 17.2 Antiemetic medicines

dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule. Oral liquid: 0.5 mg/5 mL; 2 mg/5 mL. Solid oral dosage form: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg.
metoclopramide <input checked="" type="checkbox"/>	Injection: 5 mg/mL (hydrochloride) in 2 mL ampoule. Oral liquid: 5 mg/5 mL. [c]. Tablet: 10 mg (hydrochloride). <input checked="" type="checkbox"/> Not in neonates.
<input checked="" type="checkbox"/> ondansetron <input checked="" type="checkbox"/>	Injection: 2 mg base/mL in 2 mL ampoule (as hydrochloride). Oral liquid: 4 mg base/5 mL. Solid oral dosage form: Eq 4 mg base; Eq 8 mg base; Eq 24 mg base. <input checked="" type="checkbox"/> > 1 month.

#### Complementary list

aprepitant	Capsule: 80 mg; 125 mg; 165 mg Powder for oral suspension: 125 mg in sachet
------------	--

### 17.3 Anti-Inflammatory medicines

<input checked="" type="checkbox"/> sulfasalazine	Retention enema. Suppository: 500 mg. Tablet: 500 mg.
---	---

#### Complementary List

hydrocortisone	Retention enema: 100 mg/60 mL. Suppository: 25 mg (acetate).
prednisolone	Retention enema: 20 mg/100 mL (as sodium phosphate).

### 17.4 Laxatives

<input checked="" type="checkbox"/> senna	Tablet: 7.5 mg (sennosides) (or traditional dosage forms).
Therapeutic alternatives: - bisacodyl	

### 17.5 Medicines used in diarrhoea

oral rehydration salts – zinc sulfate [c]	Co-package containing: ORS powder for dilution (see Section 17.5.1) – zinc sulfate solid oral dosage form 20 mg (see Section 17.5.2)
---	---

**WHO Model List of Essential Medicines – 23rd List (2023)****17.5.1 Oral rehydration**

oral rehydration salts	<p>Powder for dilution in 200 mL; 500 mL; 1 L.</p> <table> <tbody> <tr><td>glucose:</td><td>75 mEq</td></tr> <tr><td>sodium:</td><td>75 mEq or mmol/L</td></tr> <tr><td>chloride:</td><td>65 mEq or mmol/L</td></tr> <tr><td>potassium:</td><td>20 mEq or mmol/L</td></tr> <tr><td>citrate:</td><td>10 mmol/L</td></tr> <tr><td>osmolarity:</td><td>245 mOsm/L</td></tr> <tr><td>glucose:</td><td>13.5 g/L</td></tr> <tr><td>sodium chloride:</td><td>2.6 g/L</td></tr> <tr><td>potassium chloride:</td><td>1.5 g/L</td></tr> <tr><td>trisodium citrate dihydrate*:</td><td>2.9 g/L</td></tr> </tbody> </table> <p>*trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/L. However, as the stability of this latter formulation is very poor under tropical conditions, it is recommended only when manufactured for immediate use.</p>	glucose:	75 mEq	sodium:	75 mEq or mmol/L	chloride:	65 mEq or mmol/L	potassium:	20 mEq or mmol/L	citrate:	10 mmol/L	osmolarity:	245 mOsm/L	glucose:	13.5 g/L	sodium chloride:	2.6 g/L	potassium chloride:	1.5 g/L	trisodium citrate dihydrate*:	2.9 g/L
glucose:	75 mEq																				
sodium:	75 mEq or mmol/L																				
chloride:	65 mEq or mmol/L																				
potassium:	20 mEq or mmol/L																				
citrate:	10 mmol/L																				
osmolarity:	245 mOsm/L																				
glucose:	13.5 g/L																				
sodium chloride:	2.6 g/L																				
potassium chloride:	1.5 g/L																				
trisodium citrate dihydrate*:	2.9 g/L																				

**17.5.2 Medicines for diarrhoea**

zinc sulfate*	<p>Solid oral dosage form: 20 mg.</p> <p>*In acute diarrhoea zinc sulfate should be used as an adjunct to oral rehydration salts.</p>
---------------	---

**18. MEDICINES FOR ENDOCRINE DISORDERS****18.1 Adrenal hormones and synthetic substitutes**

hydrocortisone	Tablet: 100 micrograms (acetate).
hydrocortisone	Tablet: 5 mg; 10 mg; 20 mg.

**18.2 Androgens***Complementary List*

testosterone	Injection: 200 mg (enanthate) in 1 mL ampoule.
--------------	--

**18.3 Estrogens****18.4 Progestogens**

<input checked="" type="checkbox"/> medroxyprogesterone acetate Therapeutic alternatives: - norethisterone	Tablet: 5 mg.
--	---------------

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>18.5 Medicines for diabetes</b>	
<b>18.5.1 Insulins</b>	
insulin injection (soluble)* *including quality-assured biosimilars	Injection: 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen.
intermediate-acting insulin* *including quality-assured biosimilars	Injection: 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen (as compound insulin zinc suspension or Isophane insulin).
<input type="checkbox"/> long-acting insulin analogues* Therapeutic alternatives: - insulin degludec - insulin detemir - insulin glargine *including quality-assured biosimilars	Injection: 100 IU/mL in 3 mL cartridge or pre-filled pen.
<b>18.5.2 Oral hypoglycaemic agents</b>	
<input type="checkbox"/> empagliflozin Therapeutic alternatives: - canagliflozin - dapagliflozin	Tablet: 10 mg; 25 mg.
<input type="checkbox"/> gliclazide* Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (A10BB Sulfonylureas) metformin	Solid oral dosage form: (controlled-release tablets) 30 mg; 60 mg; 80 mg. *glibenclamide not suitable above 60 years. Tablet: 500 mg (hydrochloride).
<i>Complementary List</i>	
metformin [c]	Tablet: 500 mg (hydrochloride).
<b>18.6 Medicines for hypoglycaemia</b>	
glucagon	Injection: 1 mg/mL.
<i>Complementary List</i>	
diazoxide [c]	Oral liquid: 50 mg/mL. Tablet: 50 mg.
<b>18.7 Thyroid hormones and antithyroid medicines</b>	
levothyroxine	Tablet: 25 micrograms [c]; 50 micrograms; 100 micrograms (sodium salt).
potassium iodide	Tablet: 60 mg.
<input type="checkbox"/> methimazole Therapeutic alternatives: - carbimazole (depending on local availability)	Tablet: 5mg, 10mg, 20mg.
propylthiouracil*	Tablet: 50 mg. *For use when alternative first-line treatment is not appropriate or available; and in patients during the first trimester of pregnancy.

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>Complementary List</b>	
Lugo's solution [c]	<i>Oral liquid:</i> about 130 mg total iodine/mL.
<input type="checkbox"/> methimazole [c] Therapeutic alternatives: - carbimazole (depending on local availability)	Tablet: 5mg, 10mg, 20mg.
potassium iodide [c]	Tablet: 60 mg.
propylthiouracil* [c]	Tablet: 50 mg. <i>*For use when alternative first-line treatment is not appropriate or available</i>
<b>18.8 Medicines for disorders of the pituitary hormone system</b>	
<input type="checkbox"/> cabergoline Therapeutic alternatives: - bromocriptine	Tablet: 0.5 mg; 1 mg.
<b>Complementary List *</b>	
octreotide	<i>Injection (immediate-release):</i> 0.05 mg/mL; 0.1 mg/mL; 0.5 mg/mL (as acetate) in 1 mL vial. <i>Injection (modified-release):</i> 20 mg (as acetate) in vial plus diluent.
<b>19. IMMUNOLOGICALS</b>	
<b>19.1 Diagnostic agents</b>	
All tuberculins should comply with the WHO requirements for tuberculins.	
tuberculin, purified protein derivative (PPD)	<i>Injection.</i>
<b>19.2 Sera, Immunoglobulins and monoclonal antibodies</b>	
All plasma fractions should comply with the WHO requirements.	
anti-rabies virus monoclonal antibodies* <i>*including quality-assured biosimilars.</i>	<i>Injection:</i> 40 IU/mL in 1.25 mL, 2.5 mL vial; 100 IU/mL in 2.5 mL vial (human). <i>Injection:</i> 300 IU/mL in 10 mL vial; 600 IU/mL in 1 mL, 2.5 mL and 5 mL vial (murine).
antivenom immunoglobulin*	<i>Injection.</i> <i>*Exact type to be defined locally.</i>
diphtheria antitoxin	<i>Injection:</i> 10 000 IU; 20 000 IU in vial.
equine rabies immunoglobulin	<i>Injection:</i> 150 IU/mL; 200 IU/mL; 300 IU/mL; 400 IU/mL in vial.

**WHO Model List of Essential Medicines – 23rd List (2023)****19.3 Vaccines**

WHO Immunization policy recommendations are published in vaccine position papers based on recommendations made by the Strategic Advisory Group of Experts on Immunization (SAGE).

WHO vaccine position papers are updated three to four times per year. The list below details the vaccines for which there is a recommendation from SAGE and a corresponding WHO position paper as at March 2023. The most recent versions of the WHO position papers, reflecting the current evidence related to a specific vaccine and the related recommendations, can be accessed at any time on the WHO website at:

<https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/position-papers>

Vaccine recommendations may be universal or conditional (e.g., in certain regions, in some high-risk populations or as part of immunization programmes with certain characteristics). Details are available in the relevant position papers, and in the Summary Tables of WHO Routine Immunization Recommendations available on the WHO website at:

<https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/who-recommendations-for-routine-immunization---summary-tables>

Selection of vaccines from the Model List will need to be determined by each country after consideration of international recommendations, epidemiology and national priorities.

All vaccines should comply with the WHO requirements for biological substances.

WHO noted the need for vaccines used in children to be polyclonal.

<i>Recommendations for all</i>	
BCG vaccine	
diphtheria vaccine	
Haemophilus influenzae type b vaccine	
hepatitis B vaccine	
human papilloma virus (HPV) vaccine	
measles vaccine	
pertussis vaccine	
pneumococcal vaccine	
poliomyelitis vaccine	
rotavirus vaccine	
rubella vaccine	
tetanus vaccine	
<i>Recommendations for certain regions</i>	
Japanese encephalitis vaccine	
tick-borne encephalitis vaccine	
yellow fever vaccine	
<i>Recommendations for some high-risk populations</i>	
cholera vaccine	
dengue vaccine	
hepatitis A vaccine	
meningococcal meningitis vaccine	

## WHO Model List of Essential Medicines – 23rd List (2023)

rabies vaccine	
typhoid vaccine	
<i>Recommendations for immunization programmes with certain characteristics</i>	
influenza vaccine (seasonal)	
mumps vaccine	
varicella vaccine	
<b>20. MUSCLE RELAXANTS (PERIPHERALLY-ACTING) AND CHOLINESTERASE INHIBITORS</b>	
<input checked="" type="checkbox"/> atracurium Therapeutic alternatives to be reviewed	Injection: 10 mg/mL (besylate).
neostigmine	Injection: 500 micrograms/mL (methylsulfate) in 1 mL ampoule; 2.5 mg/mL (methylsulfate) in 1 mL ampoule. Tablet: 15 mg (bromide).
suxamethonium	Injection: 50 mg/mL (chloride) in 2 mL ampoule. Powder for Injection: (chloride), in vial.
<input checked="" type="checkbox"/> vecuronium [c] Therapeutic alternatives: -atracurium	Powder for Injection: 10 mg (bromide) in vial.
<i>Complementary List</i>	
pyridostigmine	Injection: 1 mg in 1 mL ampoule. Tablet: 60 mg (bromide).
<input checked="" type="checkbox"/> vecuronium Therapeutic alternatives to be reviewed	Powder for Injection: 10 mg (bromide) in vial.
<b>21. OPHTHALMOLOGICAL PREPARATIONS</b>	
<b>21.1 Anti-infective agents</b>	
aciclovir	Ointment: 3% w/w.
azithromycin	Solution (eye drops): 1.5%. – Trachoma
erythromycin	Ointment: 0.5% [c] – Infections due to <i>Chlamydia trachomatis</i> or <i>Neisseria gonorrhoea</i> .
<input checked="" type="checkbox"/> gentamicin Therapeutic alternatives: - amikacin - kanamycin - netilmicin - tobramycin	Solution (eye drops): 0.3% (sulfate). – <i>Bacterial blepharitis</i> – <i>Bacterial conjunctivitis</i>
natamycin	Suspension (eye drops): 5% – <i>Fungal keratitis</i>

**WHO Model List of Essential Medicines – 23rd List (2023)**

<input type="checkbox"/> ofloxacin	Solution (eye drops): 0.3%. – <i>Bacterial conjunctivitis</i> – <i>Bacterial keratitis</i>
Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01AE Fluoroquinolones)	
<input type="checkbox"/> tetracycline	Eye ointment: 1% (hydrochloride). – <i>Bacterial blepharitis</i> – <i>Bacterial conjunctivitis</i> – <i>Bacterial keratitis</i> – <i>Trachoma</i>
Therapeutic alternatives: - chlorotetracycline - oxytetracycline	
<b>21.2 Anti-inflammatory agents</b>	
<input type="checkbox"/> prednisolone	Solution (eye drops): 0.5% (sodium phosphate).
Therapeutic alternatives to be reviewed	
<b>21.3 Local anaesthetics</b>	
<input type="checkbox"/> tetracaine <input checked="" type="checkbox"/>	Solution (eye drops): 0.5% (hydrochloride). <input checked="" type="checkbox"/> Not in preterm neonates.
Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01HA Local anaesthetics) excluding cocaine and combinations	
<b>21.4 Miotics and antiglaucoma medicines</b>	
acetazolamide	Tablet: 250 mg.
latanoprost	Solution (eye drops): 50 micrograms/mL.
<input type="checkbox"/> pilocarpine	Solution (eye drops): 2%; 4% (hydrochloride or nitrate).
Therapeutic alternatives: - carbachol	
<input type="checkbox"/> timolol	Solution (eye drops): 0.25%; 0.5% (as hydrogen maleate).
Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01ED Beta blocking agents) excluding combinations	
<b>21.5 Mydriatics</b>	
<input type="checkbox"/> atropine <input checked="" type="checkbox"/>	Solution (eye drops): 0.1%; 0.5%; 1% (sulfate). <input checked="" type="checkbox"/> > 3 months.
Therapeutic alternatives: - cyclopentolate hydrochloride - homatropine hydrobromide *EMC only	
<i>Complementary List</i>	
epinephrine (adrenaline)	Solution (eye drops): 2% (as hydrochloride).
<b>21.6 Anti-vascular endothelial growth factor (VEGF) preparations</b>	
<i>Complementary List</i>	
bevacizumab*	Injection: 25 mg/mL.
*including quality-assured biosimilars	

**WHO Model List of Essential Medicines – 23rd List (2023)**

<b>22. MEDICINES FOR REPRODUCTIVE HEALTH AND PERINATAL CARE</b>	
<b>22.1 Contraceptives</b>	
<b>22.1.1 Oral hormonal contraceptives</b>	
<input type="checkbox"/> ethinylestradiol + <input type="checkbox"/> levonorgestrel Therapeutic alternatives to be reviewed	Tablet: 30 micrograms + 150 micrograms.
<input type="checkbox"/> ethinylestradiol + <input type="checkbox"/> norethisterone Therapeutic alternatives to be reviewed	Tablet: 35 micrograms + 1 mg.
levonorgestrel	Tablet: 30 micrograms; 750 micrograms (pack of two); 1.5 mg.
ulipristal	Tablet: 30 mg (as acetate).
<b>22.1.2 Injectable hormonal contraceptives</b>	
estradiol cypionate + medroxyprogesterone acetate	Injection: 5 mg + 25 mg.
medroxyprogesterone acetate	Injection (intramuscular): 150 mg/mL in 1 mL vial. Injection (subcutaneous): 104 mg/0.65 mL in pre-filled syringe or single-dose injection delivery system.
norethisterone enantate	Oily solution: 200 mg/mL in 1 mL ampoule.
<b>22.1.3 Intrauterine devices</b>	
copper-containing device	
levonorgestrel-releasing intrauterine system	Intrauterine system: with reservoir containing 52 mg of levonorgestrel
<b>22.1.4 Barrier methods</b>	
condoms	
diaphragms	
<b>22.1.5 Implantable contraceptives</b>	
etonogestrel-releasing implant	Single-rod etonogestrel-releasing implant: containing 68 mg of etonogestrel.
levonorgestrel-releasing implant	Two-rod levonorgestrel-releasing implant: each rod containing 75 mg of levonorgestrel (150 mg total).
<b>22.1.6 Intravaginal contraceptives</b>	
ethinylestradiol + etonogestrel	Vaginal ring: containing 2.7 mg + 11.7 mg
progesterone vaginal ring*	Progesterone-releasing vaginal ring: containing 2.074 g of micronized progesterone.
	*For use in women actively breastfeeding at least 4 times per day

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>22.2 Ovulation Inducers</b>	
<b>Complementary List</b>	
clomifene	Tablet: 50 mg (citrate).
<input checked="" type="checkbox"/> letrozole Therapeutic alternatives: - anastrozole	Solid oral dosage form: 2.5 mg.
<b>22.3 Uterotonics</b>	
carbetocin	Injection (heat stable): 100 micrograms/mL.
<input checked="" type="checkbox"/> ergometrine Therapeutic alternatives: - methylergometrine	Injection: 200 micrograms (hydrogen maleate) in 1 mL ampoule.
mifepristone – misoprostol  <div style="border: 1px solid black; padding: 5px;">Where permitted under national law and where culturally acceptable.</div>	Tablet 200 mg – tablet 200 micrograms.  Co-package containing: mifepristone 200 mg tablet [1] and misoprostol 200 micrograms tablet [4]  – Management of intrauterine fetal demise; – Management of induced abortion
misoprostol	Tablet: 200 micrograms. – Management of incomplete abortion and miscarriage; – Prevention and treatment of postpartum haemorrhage where oxytocin is not available or cannot be safely used  Vaginal tablet: 25 micrograms.*  *Only for use for induction of labour where appropriate facilities are available.
oxytocin	Injection: 10 IU in 1 mL.
<b>22.4 Antioxytoxics (tocolytics)</b>	
nifedipine	Immediate-release capsules: 10 mg.

**WHO Model List of Essential Medicines – 23rd List (2023)****22.5 Other medicines administered to the mother**

dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule.																														
multiple micronutrient supplement*	<p><b>Tablet containing:</b></p> <table> <tbody> <tr><td>Vitamin A (retinol acetate)</td><td>800 micrograms retinol activity equivalent</td></tr> <tr><td>Vitamin C (ascorbic acid)</td><td>70 mg</td></tr> <tr><td>Vitamin D (cholecalciferol)</td><td>5 micrograms (200 IU)</td></tr> <tr><td>Vitamin E (alpha tocopherol succinate)</td><td>10 mg alpha tocopherol equivalent</td></tr> <tr><td>Vitamin B1 (thiamine mononitrate)</td><td>1.4 mg</td></tr> <tr><td>Vitamin B2 (riboflavin)</td><td>1.4 mg</td></tr> <tr><td>Vitamin B3 (niacinamide)</td><td>18 mg niacin equivalent</td></tr> <tr><td>Vitamin B6 (pyridoxine hydrochloride)</td><td>1.9 mg</td></tr> <tr><td>Folic acid (folic acid)</td><td>680 micrograms dietary folate equivalent (400 micrograms)</td></tr> <tr><td>Vitamin B12 (cyanocobalamin)</td><td>2.6 micrograms</td></tr> <tr><td>Iron (ferrous fumarate)</td><td>30 mg</td></tr> <tr><td>Iodine (potassium iodide)</td><td>150 micrograms</td></tr> <tr><td>Zinc (zinc oxide)</td><td>15 mg</td></tr> <tr><td>Selenium (sodium selenite)</td><td>85 micrograms</td></tr> <tr><td>Copper (cupric oxide)</td><td>2 mg</td></tr> </tbody> </table> <p>*For use in specific contexts. Refer to current WHO recommendations.</p>	Vitamin A (retinol acetate)	800 micrograms retinol activity equivalent	Vitamin C (ascorbic acid)	70 mg	Vitamin D (cholecalciferol)	5 micrograms (200 IU)	Vitamin E (alpha tocopherol succinate)	10 mg alpha tocopherol equivalent	Vitamin B1 (thiamine mononitrate)	1.4 mg	Vitamin B2 (riboflavin)	1.4 mg	Vitamin B3 (niacinamide)	18 mg niacin equivalent	Vitamin B6 (pyridoxine hydrochloride)	1.9 mg	Folic acid (folic acid)	680 micrograms dietary folate equivalent (400 micrograms)	Vitamin B12 (cyanocobalamin)	2.6 micrograms	Iron (ferrous fumarate)	30 mg	Iodine (potassium iodide)	150 micrograms	Zinc (zinc oxide)	15 mg	Selenium (sodium selenite)	85 micrograms	Copper (cupric oxide)	2 mg
Vitamin A (retinol acetate)	800 micrograms retinol activity equivalent																														
Vitamin C (ascorbic acid)	70 mg																														
Vitamin D (cholecalciferol)	5 micrograms (200 IU)																														
Vitamin E (alpha tocopherol succinate)	10 mg alpha tocopherol equivalent																														
Vitamin B1 (thiamine mononitrate)	1.4 mg																														
Vitamin B2 (riboflavin)	1.4 mg																														
Vitamin B3 (niacinamide)	18 mg niacin equivalent																														
Vitamin B6 (pyridoxine hydrochloride)	1.9 mg																														
Folic acid (folic acid)	680 micrograms dietary folate equivalent (400 micrograms)																														
Vitamin B12 (cyanocobalamin)	2.6 micrograms																														
Iron (ferrous fumarate)	30 mg																														
Iodine (potassium iodide)	150 micrograms																														
Zinc (zinc oxide)	15 mg																														
Selenium (sodium selenite)	85 micrograms																														
Copper (cupric oxide)	2 mg																														
tranexamic acid	Injection: 100 mg/mL in 10 mL ampoule.																														

**22.6 Medicines administered to the neonate [c]**

caffeine citrate [c]	Injection: 20 mg/mL (equivalent to 10 mg caffeine base/mL). Oral liquid: 20 mg/mL (equivalent to 10 mg caffeine base/mL).
chlorhexidine [c]	Solution or gel: 7.1% (digluconate) delivering 4% chlorhexidine (for umbilical cord care).

**Complementary List**

<input type="checkbox"/> ibuprofen [c] Therapeutic alternatives: - indometacin	<b>Solution for injection:</b> 5 mg/mL.
<input type="checkbox"/> prostaglandin E1 [c] Therapeutic alternatives: - prostaglandin E2	<b>Solution for injection:</b> 0.5 mg/mL in alcohol.
surfactant [c]	<b>Suspension for Intratracheal Instillation:</b> 25 mg/mL or 80 mg/mL.

**23. PERITONEAL DIALYSIS SOLUTION**

<b>Complementary List</b>	
intraperitoneal dialysis solution	<b>Parenteral solution:</b> of appropriate composition.

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>24. MEDICINES FOR MENTAL AND BEHAVIOURAL DISORDERS</b>	
<b>24.1 Medicines for psychotic disorders</b>	
<input type="checkbox"/> fluphenazine Therapeutic alternatives: - haloperidol decanoate - zuclopentixol decanoate	Injection: 25 mg (decanoate or enantate) in 1 mL ampoule.
<input type="checkbox"/> haloperidol Therapeutic alternatives: - chlorpromazine	Tablet: 2 mg; 5 mg.
haloperidol	Injection: 5 mg/mL in 1 mL ampoule.
olanzapine	Powder for injection: 10 mg in vial.
<input type="checkbox"/> paliperidone Therapeutic alternatives: - risperidone injection	Injection (prolonged-release): 25 mg; 50 mg; 75 mg; 100 mg; 150 mg (as palmitate) in pre-filled syringe.
<input type="checkbox"/> risperidone Therapeutic alternatives: - aripiprazole - olanzapine - paliperidone - quetiapine	Solid oral dosage form: 0.25 mg to 6.0 mg.
<i>Complementary List</i>	
clozapine	Solid oral dosage form: 25 to 200 mg.
<b>24.2 Medicines for mood disorders</b>	
<b>24.2.1 Medicines for depressive disorders</b>	
amitriptyline	Tablet: 25 mg; 75mg (hydrochloride).
<input type="checkbox"/> fluoxetine Therapeutic alternatives: - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<b>24.2.2 Medicines for bipolar disorders</b>	
carbamazepine	Tablet (scored): 100 mg; 200 mg; 400 mg.
lithium carbonate	Solid oral dosage form: 300 mg.
<input type="checkbox"/> quetiapine Therapeutic alternatives: - aripiprazole - olanzapine - paliperidone	Tablet (immediate-release): 25 mg; 100 mg; 150 mg; 200 mg; 300 mg.  Tablet (modified-release): 50 mg; 150 mg; 200 mg; 300 mg; 400 mg.

**WHO Model List of Essential Medicines – 23rd List (2023)**

valproic acid (sodium valproate)*  *avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.	Tablet (enteric-coated): 200 mg; 500 mg.
<b>24.3 Medicines for anxiety disorders</b>	
<input type="checkbox"/> diazepam*  Therapeutic alternatives: - lorazepam	Tablet (scored): 2 mg; 5 mg.  *For short-term emergency management of acute and severe anxiety symptoms only
<input type="checkbox"/> fluoxetine  Therapeutic alternatives: - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<b>24.4 Medicines for obsessive compulsive disorders</b>	
clomipramine	Capsule: 10 mg; 25 mg (hydrochloride).
<input type="checkbox"/> fluoxetine  Therapeutic alternatives: - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<b>24.5 Medicines for disorders due to psychoactive substance use</b>	
<b>24.5.1 Medicines for alcohol use disorders</b>	
acamprosate calcium	Tablet: 333 mg
naltrexone	Injection suspension (extended-release): 380 mg in vial Tablet: 50 mg
<b>24.5.2 Medicines for nicotine use disorders</b>	
bupropion	Tablet (sustained-release): 150 mg (hydrochloride).
nicotine replacement therapy (NRT)	Chewing gum: 2 mg; 4 mg (as polacrilex). Lozenge: 2 mg; 4 mg. Oral spray: 1 mg per actuation. Transdermal patch: 5 mg to 30 mg/16 hrs; 7 mg to 21 mg/24 hrs.
varenicline	Tablet: 0.5 mg, 1 mg

**WHO Model List of Essential Medicines – 23rd List (2023)**

**24.5.3 Medicines for opioid use disorders**

**Complementary List**

methadone\* Concentrate for oral liquid: 5 mg/mL; 10 mg/mL (hydrochloride).

Therapeutic alternatives: Oral liquid: 5 mg/5 mL; 10 mg/5 mL (hydrochloride).

- buprenorphine

\*The medicines should only be used within an established support programme.

**25. MEDICINES ACTING ON THE RESPIRATORY TRACT**

**25.1 Antiflammatory medicines and medicines for chronic obstructive pulmonary disease**

<input type="checkbox"/> budesonide	Inhalation (aerosol): 100 micrograms per dose; 200 micrograms per dose.
Therapeutic alternatives: - beclometasone - ciclesonide - flunisolide - fluticasone - mometasone	
<input type="checkbox"/> budesonide + <input type="checkbox"/> formoterol	Dry powder Inhaler: 100 micrograms + 6 micrograms per dose; 200 micrograms + 6 micrograms per dose.
Therapeutic alternatives: - beclometasone + formoterol - budesonide + salmeterol - fluticasone + formoterol - fluticasone furoate + vilanterol - mometasone + formoterol	
epinephrine (adrenaline)	Injection: 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
ipratropium bromide	Inhalation (aerosol): 20 micrograms/metered dose.
<input type="checkbox"/> salbutamol	Inhalation (aerosol): 100 micrograms (as sulfate) per dose. Injection: 50 micrograms/mL (as sulfate) in 5 mL ampoule. Metered dose Inhaler (aerosol): 100 micrograms (as sulfate) per dose. Respirator solution for use in nebulizers: 5 mg/mL (as sulfate).
Therapeutic alternatives: - terbutaline	
<input type="checkbox"/> tiotropium	Powder for inhalation, capsule: 18 micrograms. Inhalation solution: 1.25 micrograms; 2.5 micrograms per actuation.
Therapeutic alternatives: - aclidinium - glycopyrronium - umeclidinium	
<b>26. SOLUTIONS CORRECTING WATER, ELECTROLYTE AND ACID-BASE DISTURBANCES</b>	
<b>26.1 Oral</b>	
oral rehydration salts	See section 17.5.1.
potassium chloride	Powder for solution.
<b>26.2 Parenteral</b>	
glucose	Injectable solution: 5% (isotonic); 10% (hypertonic); 50% (hypertonic).

## WHO Model List of Essential Medicines – 23rd List (2023)

glucose with sodium chloride	Injectable solution: 4% glucose, 0.18% sodium chloride (equivalent to Na <sup>+</sup> 30 mmol/L, Cl <sup>-</sup> 30 mmol/L). Injectable solution: 5% glucose, 0.9% sodium chloride (equivalent to Na <sup>+</sup> 150 mmol/L and Cl <sup>-</sup> 150 mmol/L); 5% glucose, 0.45% sodium chloride (equivalent to Na <sup>+</sup> 75 mmol/L and Cl <sup>-</sup> 75 mmol/L) [c].
potassium chloride	Solution: 11.2% in 20 mL ampoule (equivalent to K <sup>+</sup> 1.5 mmol/mL, Cl <sup>-</sup> 1.5 mmol/mL). Solution for dilution: 7.5% (equivalent to K 1 mmol/mL and Cl 1 mmol/mL) [c]; 15% (equivalent to K 2 mmol/mL and Cl 2 mmol/mL) [c].
sodium chloride	Injectable solution: 0.9% isotonic (equivalent to Na <sup>+</sup> 154 mmol/L, Cl <sup>-</sup> 154 mmol/L).
sodium hydrogen carbonate	Injectable solution: 1.4% isotonic (equivalent to Na <sup>+</sup> 167 mmol/L, HCO <sub>3</sub> <sup>-</sup> 167 mmol/L). Solution: 8.4% in 10 mL ampoule (equivalent to Na <sup>+</sup> 1000 mmol/L, HCO <sub>3</sub> <sup>-</sup> 1000 mmol/L).
sodium lactate, compound solution	Injectable solution.
<b>26.3 Miscellaneous</b>	
water for injection	2 mL; 5 mL; 10 mL ampoules.
<b>27. VITAMINS AND MINERALS</b>	
ascorbic acid	Tablet: 50 mg.
calcium	Tablet: 500 mg (elemental).
<input checked="" type="checkbox"/> colecalciferol [c] Therapeutic alternatives: - ergocalciferol	Oral liquid: 400 IU/mL. Solid oral dosage form: 400 IU; 1000 IU.
<input checked="" type="checkbox"/> ergocalciferol Therapeutic alternatives: - colecalciferol	Oral liquid: 250 micrograms/mL (10 000 IU/mL). Solid oral dosage form: 1.25 mg (50 000 IU).
iodine	Capsule: 190 mg. Iodized oil: 1 mL (480 mg iodine); 0.5 mL (240 mg iodine) in ampoule (oral or injectable); 0.57 mL (308 mg iodine) in dispenser bottle.
multiple micronutrient powder [c]	Sachets containing: - iron (elemental) 12.5 mg (as coated ferrous fumarate) - zinc (elemental) 5 mg - vitamin A 300 micrograms - with or without other micronutrients at recommended daily values
nicotinamide	Tablet: 50 mg.
pyridoxine	Tablet: 25 mg (hydrochloride).

**WHO Model List of Essential Medicines – 23rd List (2023)**

retinol	Capsule: 50 000 IU; 100 000 IU; 200 000 IU (as palmitate). Oral oily solution: 100 000 IU/mL (as palmitate) in multidose dispenser. Tablet (sugar-coated): 10 000 IU (as palmitate). Water-miscible injection: 100 000 IU (as palmitate) in 2 mL ampoule.
riboflavin	Tablet: 5 mg.
thiamine	Tablet: 50 mg (hydrochloride).

**Complementary List**

calcium gluconate	Injection: 100 mg/mL in 10 mL ampoule.
-------------------	--

**28. EAR, NOSE AND THROAT MEDICINES**

acetic acid [c]	Topical: 2%, in alcohol.
<input type="checkbox"/> budesonide [c]  Therapeutic alternatives to be reviewed	Nasal spray: 100 micrograms per dose.
<input type="checkbox"/> ciprofloxacin [c]  Therapeutic alternatives: - ofloxacin	Solution (ear drops): 0.3% (as hydrochloride).
<input type="checkbox"/> xylometazoline <input checked="" type="checkbox"/> [c]  Therapeutic alternatives to be reviewed	Nasal spray: 0.05%.  <input checked="" type="checkbox"/> Not in children less than 3 months.

**29. MEDICINES FOR DISEASES OF JOINTS****29.1 Medicines used to treat gout**

allopurinol	Tablet: 100 mg.
-------------	-----------------

**29.2 Disease-modifying anti-rheumatic drugs (DMARDs)**

chloroquine	Tablet: 100 mg; 150 mg (as phosphate or sulfate).
<b>Complementary List</b>	
azathioprine	Tablet: 50 mg.
hydroxychloroquine	Solid oral dosage form: 200 mg (as sulfate).
methotrexate	Tablet: 2.5 mg (as sodium).
penicillamine	Solid oral dosage form: 250 mg.
sulfasalazine	Tablet: 500 mg.

**29.3 Medicines for juvenile joint diseases****Complementary List**

acetylsalicylic acid* (acute or chronic use)	Suppository: 50 mg to 150 mg.  Tablet: 100 mg to 500 mg.
--	--

\*For use for rheumatic fever, juvenile arthritis, Kawasaki disease.

## WHO Model List of Essential Medicines – 23rd List (2023)

<b>o adalimumab*</b>	
<i>Therapeutic alternatives*:</i>	
- certolizumab pegol - etanercept - golimumab - infliximab	<b>Injection:</b> 10 mg/0.2 mL [a]; 20 mg/0.4 mL [c]; 40 mg/0.8 mL; 40 mg/0.4 mL.
*including quality-assured biosimilars	

**30. DENTAL MEDICINES AND PREPARATIONS**

fluoride	Gel: containing 2500 to 12 500 ppm fluoride (any type).  Mouthrinse: containing 230 to 900 ppm fluoride (any type).  Toothpaste, cream or gel: containing 1000 to 1500 ppm fluoride (any type).  Varnish: containing 22 500 ppm fluoride (any type).
glass ionomer cement	Single-use capsules: 0.4 g powder + 0.09 mL liquid.  Multi-use bottle: powder + liquid.  Powder (fluoro-alumino-silicate glass) contains: 25-50% silicate, 20-40% aluminium oxide, 1-20% fluoride, 15-40% metal oxide, 0-15% phosphate, remainder are polyacrylic acid powder and metals in minimal quantities. Liquid (aqueous) contains: 7-25% polybasic carboxylic acid, 45-60% polyacrylic acid.
resin-based composite (low-viscosity)*	Single-use applicator or multi-use bottle  *of any type for use as dental sealant
resin-based composite (high-viscosity)*	Single-use capsule or multi-use syringe  *of any type for use as dental filling material
silver diamine fluoride	Solution: 38% w/v.

## WHO Model List of Essential Medicines – 23rd List (2023)

Table 1.1: Medicines with age or weight restrictions

artesunate + pyronaridine tetraphosphate	> 5 kg
atropine	> 3 months
benzyl benzoate	> 2 years
betamethasone topical preparations	hydrocortisone preferred in neonates
cefazolin	> 1 month
ceftriaxone	> 41 weeks corrected gestational age
darunavir	> 3 years
dihydroartemisinin + piperazine phosphate	> 5 kg
diloxanide	> 25 kg
dolutegravir	≥ 4 weeks and ≥ 3 kg (10 mg dispersible tablet) ≥ 25 kg (50 mg tablet)
doxycycline	> 8 years (except for serious infections e.g. cholera)
ibuprofen	> 3 months (except IV form for patent ductus arteriosus)
mefloquine	> 5 kg or > 3 months
metoclopramide	Not in neonates
nevirapine	> 6 weeks
ondansetron	> 1 month
silver sulfadiazine	> 2 months
tetracaine	Not in preterm neonates
xylometazoline	> 3 months

**WHO Model List of Essential Medicines – 23rd List (2023)****Table 1.2: Explanation of dosage forms****A. Principal dosage forms used in EML – oral administration**

Term	Definition
Solid oral dosage form	<p>Refers to tablets or capsules or other solid dosage forms such as 'melts' that are immediate-release preparations. It implies that there is no difference in clinical efficacy or safety between the available dosage forms, and countries should therefore choose the form(s) to be listed depending on quality and availability.</p> <p>The term 'solid oral dosage form' is never intended to allow any type of modified-release tablet.</p>
Tablets	<p>Refers to:</p> <ul style="list-style-type: none"> <li>• uncoated or coated (film-coated or sugar-coated) tablets that are intended to be swallowed whole;</li> <li>• unscored and scored*;</li> <li>• tablets that are intended to be chewed before being swallowed;</li> <li>• tablets that are intended to be dispersed or dissolved in water or another suitable liquid before being swallowed;</li> <li>• tablets that are intended to be crushed before being swallowed.</li> </ul> <p>The term 'tablet' without qualification is never intended to allow any type of modified-release tablet.</p>
Tablets (qualified)	<p>Refers to a specific type of tablet:</p> <p><b>chewable</b> - tablets that are intended to be chewed before being swallowed;</p> <p><b>disperable</b> - tablets that are intended to be dispersed in water or another suitable liquid before being swallowed;</p> <p><b>soluble</b> - tablets that are intended to be dissolved in water or another suitable liquid before being swallowed;</p> <p><b>crushable</b> - tablets that are intended to be crushed before being swallowed;</p> <p><b>scored</b> - tablets bearing a break mark or marks where sub-division is intended in order to provide doses of less than one tablet;</p> <p><b>sublingual</b> - tablets that are intended to be placed beneath the tongue.</p> <p>The term 'tablet' is always qualified with an additional term (in parentheses) in entries where one of the following types of tablet is intended: <b>gastro-resistant</b> (such tablets may sometimes be described as enteric-coated or as delayed-release), <b>prolonged-release</b> or another modified-release form.</p>

\* Scored tablets may be divided for ease of swallowing, provided that dose is a whole number of tablets.

### WHO Model List of Essential Medicines – 23rd List (2023)

Capsules	Refers to hard or soft capsules.  The term 'capsule' without qualification is <i>never</i> intended to allow any type of modified-release capsule.
Capsules (qualified)	The term 'capsule' with qualification refers to <b>gastro-resistant</b> (such capsules may sometimes be described as enteric-coated or as delayed-release), <b>prolonged-release</b> or another modified-release form.
Granules	Preparations that are issued to patient as granules to be swallowed without further preparation, to be chewed, or to be taken in or with water or another suitable liquid.  The term 'granules' without further qualification is <i>never</i> intended to allow any type of modified-release granules.
Oral powder	Preparations that are issued to patient as powder (usually as single-dose) to be taken in or with water or another suitable liquid.
Oral liquid	Liquid preparations intended to be swallowed i.e. oral solutions, suspensions, emulsions and oral drops, including those constituted from powders or granules, but <i>not</i> those preparations intended for <i>oromucosal administration</i> e.g. gargles and mouthwashes.  Oral liquids presented as powders or granules may offer benefits in the form of better stability and lower transport costs. If more than one type of oral liquid is available on the same market (e.g. solution, suspension, granules for reconstitution), they may be interchanged and in such cases should be bioequivalent. It is preferable that oral liquids do not contain sugar and that solutions for children do not contain alcohol.

#### B. Principal dosage forms used in EML – parenteral administration

Term	Definition
Injection	Refers to solutions, suspensions and emulsions including those constituted from powders or concentrated solutions.
Injection (qualified)	Route of administration is indicated in parentheses where relevant.
Injection (oily)	The term 'injection' is qualified by '(oily)' in relevant entries.
Intravenous infusion	Refers to solutions and emulsions including those constituted from powders or concentrated solutions.

#### C. Other dosage forms

Mode of administration	Term to be used
To the eye	Eye drops, eye ointments.
Topical	For liquids: lotions, paints. For semi-solids: cream, ointment.
Rectal	Suppositories, gel or solution.
Vaginal	Pessaries or vaginal tablets.
Inhalation	Powder for inhalation, pressurized inhalation, nebulizer.

## WHO Model List of Essential Medicines – 23rd List (2023)

## Index

abacavir	19	betamethasone	42
abacavir + lamivudine	20	bevacizumab	51
abiraterone	33	bicalutamide	33
acamprosate calcium	56	bilirubin	6
acetazolamide	51	bisoprolol	37, 38, 40
acetic acid	59	bleomycin	27
acetylcysteine	4	bortezomib	32
acetylsalicylic acid	2, 26, 40, 59	budesonide	57, 59
acetylsalicylic acid + atorvastatin + ramipril	41	budesonide + formoterol	57
acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide	41	bupivacaine	1
aciclovir	19, 50	buproprion	56
adalimumab	26, 60	cabergoline	48
albendazole	7	caffeine citrate	54
alcohol based hand rub	43	calamine	42
allopurinol	34, 59	calcipotriol	42
all-trans retinoid acid (ATRA)	32	calcium	58
alteplase	40	calcium folinate (leucovorin calcium)	27
amidotriozate	43	calcium gluconate	4, 59
amikacin	9, 17	capecitabine	27
amiloride	44	carbamazepine	5, 55
amiodarone	38	carbetocin	53
amitriptyline	3, 55	carboplatin	27
amlodipine	38	cefalexin	10
amodiaquine	22	cefazolin	10
amodiaquine – sulfadoxine + pyrimethamine	24	cefiderocol	15
amoxicillin	9	cefixime	13
amoxicillin + clavulanic acid	9, 17	cefotaxime	13
amphotericin B	18, 22	ceftazidime	15
ampicillin	10	ceftazidime + avibactam	15
anastrozole	33	ceftolozane + tazobactam	15
ansuvimab	25	ceftriaxone	14
anti-D immunoglobulin	37	cefuroxime	14
anti-rabies immunoglobulin	37	charcoal, activated	4
anti-rabies virus monoclonal antibodies	48	chlorambucil	27
anti-tetanus immunoglobulin	37	chloramphenicol	10
antivenom immunoglobulin	48	chlorhexidine	43, 54
aprepitant	45	chlorine base compound	43
arsenic trioxide	27	chloroquine	23, 24, 59
artemether	22	chloroxylenol	44
artemether + lumefantrine	22	cholera vaccine	49
artesunate	23	ciclosporin	26
artesunate + amodiaquine	23	ciprofloxacin	14, 59
artesunate + mefloquine	23	cisplatin	28
artesunate + pyronaridine tetraphosphate	23	cladribine	6
ascorbic acid	58	clarithromycin	14
asparaginase	27	clindamycin	11
atazanavir + ritonavir	19	clofazimine	16, 17
atoltivimab + mabtivimab + odesvimab	25	clomifene	53
atorvastatin + perindopril + amlodipine	41	clomipramine	56
atracurium	50	clopidogrel	40
atropine	1, 4, 51	clotrimazole	18
azathioprine	26, 59	cloxacillin	11
azithromycin	13, 50	clozapine	55
barium sulfate	43	coagulation factor IX	37
BCG vaccine	49	coagulation factor VIII	37
bedaquiline	17	coal tar	42
bendamustine	27	codeine	2
benzathine benzylpenicillin	10	colecalciferol	58
benznidazole	25	colistin	15
benzoyl peroxide	42	condoms	52
benzyl benzoate	42	copper-containing device	52
benzylpenicillin	10	cryoprecipitate, pathogen-reduced	36
		cyclizine	3

## WHO Model List of Essential Medicines – 23rd List (2023)

cyclophosphamide.....	28	fentanyl.....	2
cycloserine.....	17	ferrous salt.....	35
cytarabine.....	28	ferrous salt + folic acid.....	35
dabigatran.....	35	fexinidazole.....	24
dacerbazine.....	28	filgrastim.....	33
daclatasvir.....	21	fluconazole.....	18
daclatasvir + sofosbuvir.....	21	flucytosine.....	18
dactinomycin.....	28	fludarabine.....	29
dapsone.....	16	fludrocortisone.....	46
darunavir.....	19	fluorescein.....	43
dasatinib.....	32	fluoride.....	60
daunorubicin.....	28	fluouracil.....	29, 42
deferasirox.....	36	fluoxetine.....	3, 55, 56
deferroxamine.....	4, 36	fluphenazine.....	55
delamaranid.....	17	folic acid.....	35
dengue vaccine.....	49	fomepizole.....	4
desmopressin.....	36	fosfomycin.....	15
dexamethasone.....	3, 4, 34, 45, 54	fresh-frozen plasma.....	36
dextran 70.....	37	furosemide.....	40, 44
diaphragms.....	52	gemcitabine.....	29
cliazepam.....	3, 5, 56	gentamicin.....	11, 50
diazoxide.....	47	glass ionomer cement.....	60
diethylcarbamazine.....	7	glutaramer acetate.....	6
digoxin.....	38, 40	glecaprevir + pibrentasvir.....	21
dihydroartemisinin + piperaquine phosphate.....	23	gliclazide.....	47
diltiazem.....	22	glucagon.....	47
dimercaprol.....	4	glucose.....	57
diphtheria antitoxin.....	48	glucose with sodium chloride.....	58
diphtheria vaccine.....	49	glutaral.....	44
docetaxel.....	29	glyceryl trinitrate.....	37
docusate sodium.....	3	griseofulvin.....	18
dolutegravir.....	20	Haemophilus influenzae type b vaccine.....	49
dolutegravir + lamivudine + tenofovir.....	20	haloperidol.....	3, 55
dopamine.....	40	halothane.....	1
doxorubicin.....	29	heparin sodium.....	36
doxorubicin (as pegylated liposomal).....	29	hepatitis A vaccine.....	49
doxycycline.....	11, 23, 24	hepatitis B vaccine.....	49
efavirenz.....	19	human papilloma virus (HPV) vaccine.....	49
efavirenz + emtricitabine + tenofovir.....	20	hydralazine.....	38
efavirenz + lamivudine + tenofovir.....	20	hydrochlorothiazide.....	38, 40, 44
efflomithine.....	25	hydrocortisone.....	4, 34, 42, 45, 46
empagliflozin.....	47	hydroxocobalamin.....	35
emtricitabine + tenofovir.....	20	hydroxy-carbamide (hydroxyurea).....	30, 36
enalapril.....	38, 40	hydroxychloroquine.....	59
enoxaparin.....	36	hyoscine butylbromide.....	3
entecavir.....	21	hyoscine hydrobromide.....	3
ephedrine.....	1	ibrutinib.....	32
epinephrine (adrenaline).....	4, 38, 51, 57	Ibuprofen.....	2, 26, 54
equine rabies immunoglobulin.....	48	ifosfamide.....	30
ergocaliferol.....	58	imatinib.....	32
ergometrine.....	53	influenza vaccine.....	50
erlotinib.....	32	insulin injection (soluble).....	47
erythromycin.....	50	intermediate-acting insulin.....	47
erythropoiesis-stimulating agents.....	35	intraperitoneal dialysis solution.....	54
estradiol cypionate + medroxyprogesterone acetate .....	52	iodine.....	58
ethambutol.....	16	iohexol.....	43
ethambutol + Isoniazid + pyrazinamide + rifampicin .....	16	ipratropium bromide.....	57
ethambutol + Isoniazid + rifampicin .....	16	Irinotecan.....	30
ethanol.....	43	isoflurane.....	1
ethinylestradiol + etonogestrel.....	52	isoniazid.....	16
ethinylestradiol + levonorgestrel.....	52	isoniazid + pyrazinamide + rifampicin .....	16
ethinylestradiol + norethisterone .....	52	Isoniazid + pyridoxine + sulfamethoxazole + trimethoprim.....	20
ethionamide.....	16, 17	isoniazid + rifampicin .....	16
ethosuximide .....	6	Isoniazid + rifapentine .....	16
etonogestrel-releasing implant .....	52	isosorbide dinitrate .....	37
etoposide .....	29	itraconazole .....	18
everolimus .....	32	ivermectin .....	7, 25

## WHO Model List of Essential Medicines – 23rd List (2023)

Japanese encephalitis vaccine.....	49
ketamine .....	1
lactulose.....	3
lamivudine.....	19
lamivudine + zidovudine.....	20
lamotrigine .....	5
latanoprost .....	51
ledipasvir + sofosbuvir .....	21
<i>lenalidomide</i> .....	33
<i>letrozole</i> .....	53
<i>leuprorelin</i> .....	34
levamisole .....	7
levetiracetam.....	5, 6
levodopa + carbidopa .....	6
<i>levofloxacin</i> .....	17
levonorgestrel .....	52
levonorgestrel-releasing implant.....	52
levonorgestrel-releasing intrauterine system.....	52
levothyroxine .....	47
lidocaine.....	1, 38
lidocaine + epinephrine (adrenaline).....	1
<i>linezolid</i> .....	15, 17
lisinopril + amlodipine .....	39
lisinopril + hydrochlorothiazide .....	39
lithium carbonate .....	55
long-acting insulin analogues .....	47
loperamide .....	3
lopinavir + ritonavir .....	19
loratadine .....	4
lorazepam .....	5
losartan .....	39, 40
<i>Lugo's solution</i> .....	48
magnesium sulfate.....	5
mannitol .....	44
measles vaccine .....	49
mebendazole .....	7
medroxyprogesterone acetate .....	46, 52
mefloquine .....	23, 24
meglumine antimoniate .....	22
<i>meglumine lotoxate</i> .....	43
melarsoprol .....	25
<i>melphalan</i> .....	30
meningococcal meningitis vaccine .....	49
<i>mercaptopurine</i> .....	30
meropenem .....	15, 17
meropenem + vaborbactam .....	16
mesna .....	34
metformin .....	47
<i>methadone</i> .....	2, 57
methylazole .....	47, 48
<i>methotrexate</i> .....	30, 42, 59
methyldopa .....	39
<i>methylprednisolone</i> .....	34
methylene blue (methylthioninium chloride) .....	4
metoclopramide .....	3, 45
metronidazole .....	12, 22
<i>microfungin</i> .....	18
micronazole .....	41
midazolam .....	1, 3, 5
misopristone – mlsoprostol .....	53
mitfenesine .....	22
misoprostol .....	53
morphine .....	1, 2
moxifloxacin .....	16, 17
multiple micronutrient powder .....	58
multiple micronutrient supplement .....	54
mumps vaccine .....	50
mupirocin .....	42
naloxone .....	4
naltrexone .....	56
<i>nizatmycin</i> .....	50
neostigmine .....	50
nevirapine .....	19
niclosamide .....	7
nicotinamide .....	58
nicotine replacement therapy (NRT) .....	56
nifedipine .....	53
nifurtimox .....	25
<i>nitotinib</i> .....	32
nitrofurantoin .....	12
nitrous oxide .....	1
<i>nivolumab</i> .....	33
norethisterone enantate .....	52
<i>normal immunoglobulin</i> .....	37
nystatin .....	18
octreotide .....	48
ofloxacin .....	51
olanzapine .....	55
omeprazole .....	44
ondansetron .....	3, 45
oral rehydration salts .....	46, 57
oral rehydration salts – zinc sulfate .....	45
<i>oseltamivir</i> .....	21
<i>oxaliplatin</i> .....	30
<i>oxamniquine</i> .....	7
oxygen .....	1, 2
oxytocin .....	53
<i>paclitaxel</i> .....	31
paliperidone .....	55
<i>p-aminosalicylate sodium</i> .....	17
<i>pancreatic enzymes</i> .....	44
paracetamol (acetaminophen) .....	2, 26
paromomycin .....	22
<i>pegaspargase</i> .....	31
<i>pegfilgrastim</i> .....	33
penicillamine .....	4, 59
<i>pentamidine</i> .....	24
permethrin .....	42
pertussis vaccine .....	49
phenobarbital .....	5
phenoxymethylpenicillin .....	12
phenytoin .....	5
phytomenadione .....	36
pilocarpine .....	51
piperacillin + tazobactam .....	15
platelets .....	36
<i>plazomicin</i> .....	16
pneumococcal vaccine .....	49
podophyllum resin .....	42
poliomylitis vaccine .....	49
<i>polymyxin B</i> .....	16
potassium chloride .....	57, 58
potassium ferric hexacyano-ferrate(II) -2H <sub>2</sub> O .....	4
potassium iodide .....	18, 47, 48
potassium permanganate .....	42
povidone iodine .....	43
praziquantel .....	7
prednisolone .....	4, 34, 45, 51
<i>pretermind</i> .....	17
primaquine .....	23
procaine benzylpenicillin .....	12
<i>procarbazine</i> .....	31
progesterone vaginal ring .....	52
proguanil .....	24

**WHO Model List of Essential Medicines – 23rd List (2023)**

propofol	1	succimer	4
propranolol	26	sulfadiazine	24
propythiouracil	47, 48	sulfadoxine + pyrimethamine	23, 24
prostaglandin E1	54	sulfamethoxazole + trimethoprim	12, 24
protamine sulfate	36	sulfasalazine	45, 59
pyrantel	7	sumatriptan	26
pyrazinamide	16	suramin sodium	24
pyridostigmine	50	surfactant	54
pyridoxine	58	suxamethonium	50
pyrimethamine	24	tacrolimus	26
quetiapine	55	tamoxifen	34
quinine	23	telmisartan + amlodipine	39
rabies vaccine	50	telmisartan + hydrochlorothiazide	39
raltegravir	20	tenofovir disoproxil fumarate	19, 21
ranitidine	44	terbinafine	41
rasburicase	35	testosterone	46
ravidasvir	21	tetanus vaccine	49
ready-to-use therapeutic food	35	tetracaine	51
realgar-Indigo naturals formulation	31	tetracycline	51
red blood cells	36	thalidomide	33
resin-based composite (high-viscosity)	60	thiamine	59
resin-based composite (low-viscosity)	60	tick-borne encephalitis vaccine	49
retinol	59	timolol	51
ribavirin	20, 21	tioguanine	31
riboflavin	59	tiotropium	57
rifabutin	16	tranexameric acid	36, 54
rifampicin	16	trastuzumab	32
rifapentine	16	triamicinolone hexacetonide	60
risperidone	55	triclabendazole	7
ritonavir	19	trimethoprim	13
rituximab	6, 32	tropicamide	43
rotavirus vaccine	49	tuberculin, purified protein derivative (PPD)	48
rubella vaccine	49	typhoid vaccine	50
salbutamol	57	ulipristal	52
salicylic acid	42	urea	42
seelenium sulfide	41	valganciclovir	20, 21
senna	3, 45	valproic acid (sodium valproate)	5, 6, 56
sevoflurane	1	vancocin	15
silver diamine fluoride	60	varenicline	56
silver sulfadiazine	42	varicella vaccine	50
simvastatin	41	vecuronium	50
sodium calcium edetate	4	verapamil	37, 38
sodium chloride	58	vinblastine	31
sodium hydrogen carbonate	58	vincristine	31
sodium lactate	58	vinorelbine	31
sodium nitrite	4	voriconazole	18
sodium nitroprusside	39	warfarin	36
sodium stibogluconate	22	water for injection	58
sodium thiosulfate	4, 41	whole blood	37
sofosbuvir	21	xylometazoline	59
sofosbuvir + velpatasvir	21	yellow fever vaccine	49
spectinomycin	12	zidovudine	19
spironolactone	40, 44	zinc sulfate	46
streptokinase	40	zoledronic acid	35
streptomycin	17		

[No. F. 8-43/2021-DD(PS).]

**SIBGHATULLAH,**  
**Section Officer (Admn-II).**

